arXiv:1102.5055v1 [math.FA] 24 Feb 2011

Maximal monotone operators with non-empty domain interior; characterizations and continuity properties

M.D. Voisei

Abstract

In the context of general Banach spaces characterizations for the maximal monotonicity of operators with non-empty domain interior as well as stronger continuity properties for such operators are provided.

1 Introduction

In the context of locally convex or Banach spaces there are few characterizations for the maximal monotonicity of an operator (see e.g. [10, Theorem 3.8], [22, Theorem 2.3], and [26, Theorem 6]). All these characterizations are based on special convex representations associated to the operator.

In a finite-dimensional space a complete characterization for the maximality of a monotone operator is given in [13, Theorem 3.4] in terms of direct operator notions: the near convexity of its domain, convexity of its values, graph closedness, and behavior at the boundary of its domain.

In a Banach space context characterizations of maximality, similar to those found in the finite-dimensional case, are available for full-space or open convex domain monotone multi-functions (see [4, Theorem 1.2], [8, Lemma 2.2], [21, Theorem 40.2, p. 155], [24, Lemma 4.2], and [17, Lemma 7.7, p. 104]).

Every maximal monotone operator in a finite dimensional space has a non-empty convex relative interior of its domain which is dense in the domain and a convex domain closure (see e.g. [15], [19, Theorems 6.2, 6.3], [20, Theorem 12.41, p. 554]).

That is why, characterizations of the maximality of a monotone operator with non-empty relative (algebraic) domain interior in a general Banach space, in terms of notions directly linked to the operator and similar to those seen in the finite-dimensional case, constitute generalizations of all fore-mentioned results and that is our primary goal.

Our secondary goal is to reveal several continuity properties with respect to the strong×weak-star topology for maximal monotone operators which have non-empty domain interiors and are defined in a normed barrelled space, such as, the closedness of the graph, the upper semicontinuity, and the Cesari property.

The plan of the paper is as follows. The next section introduces the reader to the main notions and notations used in this article. Section 3 dwells with the restrictions of an operator to affine subsets. Section 4 analyzes the finite-dimensional case and provides a new proof of [13, Theorem 3.4]. In Section 5 the characterizations previously seen in the finite dimensional context are extended to arbitrary Banach spaces via a hemicontinuity condition, demiclosedness, or representability. Section 6 deals with the continuity properties of monotone demiclosed operators that have a non-empty domain interior.

2 Preliminaries

Let (E,μ) be a locally convex space and $A \subset E$. We denote by "$\operatorname{conv} A$" the *convex hull* of A, "$\operatorname{aff} A$" the *affine hull* of A, "$\operatorname{lin} A$" the *linear hull* of A; "$\operatorname{cl}_\mu(A) = \overline{A}^{\mu}$" the $\mu-$*closure* of A, "$\operatorname{ri}_\mu A$" the topological interior of A with respect to $\operatorname{cl}_\mu(\operatorname{aff} A)$, "$\operatorname{core} A$" the *algebraic interior* of A, "${}^{i}A$" the *relative algebraic interior* of A, and ${}^{\mu-ic}A := {}^{i}A$ if $\operatorname{aff} A$ is $\mu-$closed and ${}^{\mu-ic}A := \emptyset$ otherwise, the relative algebraic interior of A with respect to $\operatorname{cl}_\mu(\operatorname{aff} A)$.

When the topology μ is implicitly understood (such is the case when we deal with the strong topology of a normed space) the use of the $\mu-$notation is avoided.

For $f, g : E \to \overline{\mathbb{R}} := \mathbb{R} \cup \{-\infty, +\infty\}$ we set $[f \le g] := \{x \in E \mid f(x) \le g(x)\}$; the sets $[f = g]$, $[f < g]$, and $[f > g]$ are defined similarly.

Throughout this paper, if not otherwise explicitly mentioned, $(X, \|\cdot\|)$ is a non trivial (that is, $X \neq \{0\}$) normed space, X^* is its topological dual endowed with the weak-star topology w^*, the topological dual of (X^*, w^*) is identified with X, the weak topology on X is denoted by w, and the strong topology on X is denoted by s. The closed unit ball of X is denoted by $B_X := \{x \in X \mid \|x\| \le 1\}$. The *duality product* of $X \times X^*$ is denoted by $\langle x, x^*\rangle := x^*(x) =: c(x, x^*)$, for $x \in X$, $x^* \in X^*$.

As usual, for $S \subset X$, $S^\perp := \{x^* \in X^* \mid \langle x, x^*\rangle = 0$, for every $x \in S\}$, $\sigma_S(x^*) := \sup_{x \in S}\langle x, x^*\rangle$, $x^* \in X^*$ and for $A \subset X^*$, $A^\perp := \{x \in X \mid \langle x, x^*\rangle = 0$, for every $x \in A\}$, $\sigma_A(x) = \sup_{x^* \in A}\langle x, x^*\rangle$, $x \in X$.

To a multifunction $T : X \rightrightarrows X^*$ we associate its *graph:* $\operatorname{Graph} T = \{(x, x^*) \in X \times X^* \mid x^* \in T(x)\}$, *inverse:* $T^{-1} : X^* \rightrightarrows X$, $\operatorname{Graph} T^{-1} = \{(x^*, x) \mid (x, x^*) \in \operatorname{Graph} T\}$, *domain*: $D(T) := \{x \in X \mid T(x) \neq \emptyset\} = \operatorname{Pr}_X(\operatorname{Graph} T)$, and *range*: $R(T) := \{x^* \in X^* \mid x^* \in T(x)$ for some $x \in X\} = \operatorname{Pr}_{X^*}(\operatorname{Graph} T)$. Here Pr_X and Pr_{X^*} are the projections of $X \times X^*$ onto X and X^*, respectively. When no confusion can occur, T will be identified with $\operatorname{Graph} T$.

On X, we consider the following classes of functions and operators

$\Lambda(X)$ the class formed by proper convex functions $f : X \to \overline{\mathbb{R}}$. Recall that f is *proper* if $\operatorname{dom} f := \{x \in X \mid f(x) < \infty\}$ is nonempty and f does not take the value $-\infty$,

$\Gamma_\tau(X)$ the class of functions $f \in \Lambda(X)$ that are τ–lower semicontinuous (τ–lsc for short); when the topology is implicitly understood we use the notation $\Gamma(X)$,

$\mathcal{M}(X)$ the class of non-empty monotone operators $T : X \rightrightarrows X^*$. Recall that $T : X \rightrightarrows X^*$ is *monotone* if $\langle x_1 - x_2, x_1^* - x_2^* \rangle \geq 0$, for all $(x_2, x_2^*), (x_2, x_2^*) \in T$.

$\mathscr{M}(X)$ the class of maximal monotone operators $T : X \rightrightarrows X^*$. The maximality is understood in the sense of graph inclusion as subsets of $X \times X^*$.

To a proper function $f : X \to \overline{\mathbb{R}}$ and a topology τ on X we associate:

- the *epigraph* of f: $\operatorname{epi} f := \{(x,t) \in X \times \mathbb{R} \mid f(x) \leq t\}$,
- the *convex hull* of f: $\operatorname{conv} f : X \to \overline{\mathbb{R}}$, is the greatest convex function majorized by f, $(\operatorname{conv} f)(x) := \inf\{t \in \mathbb{R} \mid (x,t) \in \operatorname{conv}(\operatorname{epi} f)\}$, $x \in X$,
- the *$\tau-$lsc convex hull* of f: $\operatorname{cl}_\tau \operatorname{conv} f : X \to \overline{\mathbb{R}}$, is the greatest τ–lsc convex function majorized by f, $(\operatorname{cl}_\tau \operatorname{conv} f)(x) := \inf\{t \in \mathbb{R} \mid (x,t) \in \operatorname{cl}_\tau \operatorname{conv} \operatorname{epi} f\}$, $x \in X$,
- the *convex conjugate* of $f : X \to \overline{\mathbb{R}}$ with respect to the dual system (X, X^*): $f^* : X^* \to \overline{\mathbb{R}}$, $f^*(x^*) := \sup\{\langle x, x^* \rangle - f(x) \mid x \in X\}$, $x^* \in X^*$.
- the *subdifferential* of f at $x \in X$: $\partial f(x) := \{x^* \in X^* \mid \langle x' - x, x^* \rangle + f(x) \leq f(x'),\ \forall x' \in X\}$ for $x \in \operatorname{dom} f$; $\partial f(x) := \emptyset$ for $x \notin \operatorname{dom} f$. Recall that $N_C = \partial I_C$ is the *normal cone* of $C \subset X$, where I_C is the *indicator function* of $C \subset X$ defined by $I_C(x) := 0$ for $x \in C$ and $I_C(x) := \infty$ for $x \in X \setminus C$.

Let $Z := X \times X^*$. It is known that $(Z, s \times w^*)^* = Z$ via the coupling

$$z \cdot z' := \langle x, x'^* \rangle + \langle x', x^* \rangle\,, \quad \text{for } z = (x, x^*),\ z' = (x', x'^*) \in Z.$$

For a proper function $f : Z \to \overline{\mathbb{R}}$ all the above notions are defined similarly. The conjugate of f with respect to the natural dual system (Z, Z) induced by the previous coupling is given by

$$f^\square : Z \to \overline{\mathbb{R}}, \quad f^\square(z) = \sup\{z \cdot z' - f(z') \mid z' \in Z\},$$

and by the biconjugate formula, $f^{\square\square} = \operatorname{cl}_{s \times w^*} \operatorname{conv} f$ whenever $f^\square$ or $\operatorname{cl}_{s \times w^*} \operatorname{conv} f$ is proper.

We consider the following classes of functions on Z:

$$\begin{aligned} \mathcal{C} :&= \mathcal{C}(Z) := \{f \in \Lambda(Z) \mid f \geq c\}, \\ \mathcal{R} :&= \mathcal{R}(Z) := \Gamma_{s \times w^*}(Z) \cap \mathcal{C}(Z), \\ \mathcal{D} :&= \mathcal{D}(Z) := \{f \in \mathcal{R}(Z) \mid f^\square \geq c\}. \end{aligned}$$

It is known that $[f = c] \in \mathcal{M}(X)$ for every $f \in \mathcal{C}(Z)$ (see e.g. [24, Lemma 3.1]).

To a multi-valued operator $T : X \rightrightarrows X^*$ we associate the following functions:

- the *Fitzpatrick function* of T (introduced in [10]): $\varphi_T : Z \to \overline{\mathbb{R}}$, $\varphi_T := c_T^{\square}$, where $c_T : Z \to \overline{\mathbb{R}}$, $c_T := c + \iota_T$;
- $\psi_T := \operatorname{cl}_{s\times w^*} \operatorname{conv} c_T$ (first considered in [22]); $\psi_T = \varphi_T^{\square} = c_T^{\square\square}$ whenever φ_T or ψ_T is proper (for example when $T \in \mathcal{M}(X)$ (see e.g. [23, Proposition 3.2])).

Note that for every $T \subset Z$, $[\varphi_T \le c]$ describes the set of all $z \in Z$ that are monotonically related (m.r. for short) to T, that is, $c(z - a) \ge 0$, for every $a \in T$.

We call a multifunction $T : X \rightrightarrows X^*$

unique if T admits a unique maximal monotone extension;

representable in Z if $T = [f = c]$, for some $f \in \mathcal{R}$; in this case f is called a *representative* of T. We denote by $\mathcal{R}_T$ the class of representatives of T; this notion was first considered in this form in [22];

dual-representable if $T = [f = c]$, for some $f \in \mathcal{D}$; in this case f is called a *d–representative* of T and we denote by $\mathcal{D}_T$ the class of d-representatives of T;

NI or of *negative infimum type in Z* if $\varphi_T \ge c$ in Z;

demiclosed if $\operatorname{Graph} T$ is closed with respect to the strong×weak-star convergence of bounded nets in Z, that is, if $x_i \to x_0$ strongly in X, $x_i^* \to x_0^*$ weakly-star in X^*, $(x_i^*)_i$ is (strongly) bounded in X^*, and $\{(x_i, x_i^*)\}_i \subset T$ then $(x_0, x_0^*) \in T$.

$strongly \times weakly - star\ upper\ \mathscr{H} - semicontinuous$ ($s \times w^*-$usc for short) *at* $x \in X$ if for every weak-star open set $V \supset Tx$ there exists a neighborhood U of x such that $T(U) := \cup_{u\in U} Tu \subset V$ (or equivalently for every net $\{x_i\}_{i\in I}$ with $x_i \to x$, strongly in X, eventually $Tx_i \subset V$). The operator T is $s \times w^*-usc$ if T is $s \times w^*-$usc at x, for every $x \in X$.

It is easily checked that every representable operator is monotone demiclosed and has w^*-closed convex values.

Recall that $T \in \mathscr{M}(X)$ iff T is NI and representable (see [22] or [24]); also, whenever $T \in \mathscr{M}(X)$

T is dual-representable with $\varphi_T, \psi_T \in \mathcal{D}_T$ (see [22, Theorems 2.1, 2.2]),

$\operatorname{ri} D(T) = {}^{\mathrm{ic}}D(T) = {}^{\mathrm{ic}}(\operatorname{ri} D(T))$, in particular $\operatorname{int} D(T) = \operatorname{core} D(T)$ and $\overline{D(T)}$ is convex whenever ${}^{\mathrm{ic}}(\operatorname{conv} D(T)) \neq \emptyset$ (see [27, Corollary 3]).

Some of the main characterizations of maximal monotonicity can be found in [10, Theorem 3.8], [22, Theorem 2.3], [26, Theorem 6], [24, Lemma 4.2], [21, Theorem 40.2, p. 155], [16, Lemma 7.7, p. 104] and they will often be recalled in the sequel. For other properties of the notions discussed in this section we suggest [26, 27, 23, 24, 22, 25].

Since the characterization of maximality for a monotone operator with a singleton domain is trivial, in this paper we do not consider singleton-domain operators.

Throughout this article the conventions $\sup\emptyset = -\infty$ and $\inf\emptyset = \infty$ are enforced.

3 Restrictions to affine sets

For X a separated locally convex space and $F \subset X$ a linear subspace with $D(T)\cap F \neq \emptyset$, let $T_F := \iota_F^* T \iota_F : F \rightrightarrows F^*$

$$T_F x := \{x^*|_F \mid x^* \in Tx\},\ x \in D(T_F) := D(T)\cap F. \tag{1}$$

Here $\iota_F^* : X^* \to F^*$, $\iota_F^*(x^*) = x^*|_F$ stands for the adjoint of $\iota_F : F \to X$, $\iota_F(x) = x$.

Lemma 1 *Let X be a normed space, let $T : X \rightrightarrows X^*$, and let $F \subset X$ be a linear subspace such that $D(T)\cap F \neq \emptyset$.*

Consider the conditions: (i) $T_F \in \mathscr{M}(F)$; (ii) $T + N_F \in \mathscr{M}(X)$.

Then (ii) ⇒ (i). If, in addition, F is closed then (i) ⇒ (ii).

Proof. It is easily checked that $T_F \in \mathcal{M}(F)$ iff $T|_F \in \mathcal{M}(X)$ iff $T + N_F \in \mathcal{M}(X)$. Here $T|_F$ stands for the restriction of T to F.

(ii) ⇒ (i) Let $(f, f^*) \in F\times F^*$ be m.r. to T_F, i.e.,

$$\langle f - x, f^* - x^*|_F\rangle \geq 0,\ \forall x \in D(T)\cap F,\ x^* \in Tx. \tag{2}$$

Take $y^* \in X^*$ with $y^*|_F = f^*$. Relation (2) implies that (f, y^*) is m.r. to $T + N_F \in \mathscr{M}(X)$; from which $f \in D(T)$ and $y^* \in Tf + F^\perp$. This yields $(f, f^*) \in T_F$.

(i) ⇒ (ii) Assume that F is closed and $T_F \in \mathscr{M}(F)$. Let (x_0, x_0^*) be m.r. to $T + N_F$, that is,

$$\langle x_0 - x, x_0^* - x^* - f^*\rangle \geq 0,\ \forall x \in D(T)\cap F,\ x^* \in Tx,\ f^* \in N_F(x) = F^\perp. \tag{3}$$

Since $F^\perp$ is a linear subspace, $x_0 - x \in F^{\perp\perp} = F$ for every $x \in D(T)\cap F$; in particular $x_0 \in F$. Relation (3) becomes $\langle x_0 - x, x_0^* - x^*\rangle \geq 0$, for every $x \in D(T)\cap F$, $x^* \in Tx$. In other words $(x_0, x_0^*|_F)$ is m.r. to T_F and so $x_0 \in D(T)$ and there is $x_1^* \in Tx_0$ such that $x_0^*|_F = x_1^*|_F$. This last equality is equivalent to $x_0^* - x_1^* \in F^\perp = N_F(x_0)$. ■

Remark 1 *Notice that for $T = X\times\{0\}$, $T_F = F\times\{0\} \in \mathscr{M}(F)$, for every linear subspace $F \subset X$, while $T + N_F = F \times F^\perp \in \mathscr{M}(X)$ iff F is closed. Therefore a closed F is necessary and sufficient for (i) ⇒ (ii) to hold.*

Lemma 2 *Let X be a normed space, let $T : X \rightrightarrows X^*$, and let F be a linear subspace such that $D(T) \subset F$.*

Consider the conditions: (i) $T \in \mathscr{M}(X)$; (ii) $T_F \in \mathscr{M}(F)$ *and* $T = T + N_F$.

Then (i) ⇒ (ii). If, in addition, F is closed then (ii) ⇒ (i).

Proof. (i) $\Rightarrow$ (ii) Since $\mathscr{M}(X) \ni T \subset T + N_F \in \mathcal{M}(X)$ we find $T = T + N_F \in \mathscr{M}(X)$. Hence, from Lemma 1, $T_F \in \mathscr{M}(F)$.
(ii) $\Rightarrow$ (i) According to Lemma 1, $T = T + N_F \in \mathscr{M}(X)$ since $T_F \in \mathscr{M}(F)$. $\blacksquare$

Remark 2 *For a linear subspace $F \subset X$, take $T = N_F = F \times F^{\perp}$. Then $T_F = F \times \{0\} \in \mathscr{M}(F)$, $T = T + N_F$, but $T \in \mathscr{M}(X)$ iff F is closed. Therefore, in the previous lemma, the implication (ii) $\Rightarrow$ (i) requires F to be closed.*

For $T : X \rightrightarrows X^*$ and $z \in X$, consider the translation $T_z(x) := T(x + z)$, $x \in D(T_z) = D(T) - z$. Similarly, for an affine subset $A \subset X$ and $z \in A$ let $F := A - z$ be the linear subspace parallel to A. Note that $\operatorname{Graph} N_A = A \times F^{\perp}$. By definition $T_{A,z} : F \rightrightarrows F^*$, $T_{A,z} := (T_z)_F$ or equivalently

$$\operatorname{Graph} T_{A,z} = \{(x, x^*|_F) \mid x^* \in T(z+x),\ x \in D(T_{A,z}) := (D(T) - z) \cap F\}. \quad (4)$$

Notice that $T_{A,z}$ is non-empty iff $D(T) \cap A \neq \emptyset$. Also, note that $T_{F,z} = (T_F)_z$, for every $z \in F$.

The following two results are consequences of the previous lemmas and the fact that a translation preserves the NI type and the (maximal) monotonicity; in other words T is NI iff T_z is NI and T is (maximal) monotone iff T_z is (maximal) monotone, for every (some) $z \in X$.

Lemma 3 *Let X be a normed space, let $T : X \rightrightarrows X^*$, let $A \subset X$ be an affine set such that $D(T) \cap A \neq \emptyset$, and let F be the linear subspace parallel to A. If $T + N_A \in \mathscr{M}(X)$ then $T_{A,z} \in \mathscr{M}(F)$, for every $z \in A$. If, in addition, A is closed then $T + N_A \in \mathscr{M}(X)$ whenever $T_{A,z} \in \mathscr{M}(F)$, for some $z \in A$.*

Proof. If $T + N_A \in \mathscr{M}(X)$ then $(T + N_A)_z = T_z + N_F \in \mathscr{M}(X)$. According to Lemma 1, $T_{A,z} \in \mathscr{M}(F)$ for every $z \in A$.

Conversely, if A is closed and $T_{A,z} = (T_z)_F \in \mathscr{M}(F)$ for some $z \in A$ then, from Lemma 1, $T_z + N_F = (T + N_A)_z, T + N_A \in \mathscr{M}(X)$. $\blacksquare$

Similarly, one has

Lemma 4 *Let X be a normed space, let $T : X \rightrightarrows X^*$, let $A \subset X$ be an affine set such that $D(T) \subset A$, and let F be the linear subspace parallel to A. If $T \in \mathscr{M}(X)$ then $T = T + N_A$ and $T_{A,z} \in \mathscr{M}(F)$, for every $z \in A$. If, in addition, A is closed then $T \in \mathscr{M}(X)$ whenever $T = T + N_A$ and $T_{A,z} \in \mathscr{M}(F)$, for some $z \in A$.*

Hereditary properties from T to $T_{A,z}$, where A is affine with $D(T) \subset A$, are studied next.

Proposition 5 *Let X be a separated locally convex space, let $T : X \rightrightarrows X^*$ be such that $T = T + N_A$, where $A \subset X$ is affine with $D(T) \subset A$, and let F be the linear subspace parallel to A. Then T has w^*-closed values in X^* whenever $T_{A,z}$ has w^*-closed values in F^*, for some $z \in A$. If, in addition, A is closed then $T_{A,z}$ has w^*-closed values in F^*, for every $z \in A$ whenever T has w^*-closed values in X^*.*

Proof. We may assume without loss of generality that $z = 0 \in A = F$ in which case $T_{A,z} = T_F$.

It suffices to observe that $(\iota_F^*)^{-1}(T_F x) = Tx + F^\perp = Tx$, for every $x \in D(T)$ to conclude that T has w^*-closed values in X^* provided that T_F has w^*-closed values in F^*.

Conversely, assume that F is closed. Let $\pi : (X^*, w^*) \to (X^*, w^*)/F^\perp$, $\pi(x^*) = x^* + F^\perp$, $x^* \in X^*$, be the projection map of (X^*, w^*) onto the quotient space $(X^*, w^*)/F^\perp$. Note that $\pi^{-1}(\pi(Tx)) = Tx + F^\perp = Tx$ is w^*-closed convex, i.e., $\pi(Tx)$ is closed convex in $(X^*, w^*)/F^\perp$, for every $x \in D(T)$.

Since F is closed in X the locally convex spaces $(X^*, w^*)/F^\perp$ and (F^*, w^*) are isomorphic via $j : (X^*, w^*)/F^\perp \to (F^*, w^*)$, $j(x^* + F^\perp) = x^*|_F$, $x^* \in X^*$. Notice that $T_F x = \iota_F^*(Tx) = j(\pi(Tx))$, $x \in D(T)$, to conclude that T_F has w^*-closed convex values. ■

Remark 3 *If T has w^*-closed values in X^* but F is not closed then one cannot expect T_F to have w^*-closed values in F^* even though $T = T + N_F$. Indeed, let F be a proper dense linear subspace of a separated locally convex space X. Then $\iota_F^* : (X^*, w^*) \to (F^*, w^*)$ is a continuous bijection but not an isomorphism, since $F \subsetneq X$. Take S a w^*-closed subset of X^* such that $\iota_F^*(S)$ is not w^*-closed in F^* and $T := \{0\} \times S \subset X \times X^*$. Then $T_F = \{0\} \times \iota_F^*(S)$ does not have w^*-closed values (and implicitly is not $s \times w^*-$closed in $F \times F^*$) while T is $s \times w^*-$closed in $X \times X^*$, has w^*-closed values, and $T = T + N_F$, since $N_F = F \times \{0\}$.*

Remark 4 *Let $T : X \rightrightarrows X^*$ and let $A \subset X$ be affine with $D(T) \subset A$. It is trivial to check that $T_{A,z}$ has convex values, for every $z \in A$ whenever T has convex values. Conversely, T has convex values whenever $T_{A,z}$ has convex values, for some $z \in A$, provided, in addition, that $T = T + N_A$.*

Proposition 6 *Let X be a separated locally convex space, let $T : X \rightrightarrows X^*$ be such that $T = T + N_A$, where $A \subset X$ is an affine set such that $D(T) \subset A$, and let F be the linear subspace parallel to A. Then T is $s \times w^*-$closed in $X \times X^*$ whenever $T_{A,z}$ is $s \times w^*-$closed in $F \times F^*$, for some $z \in A$. If, in addition, A is closed then $T_{A,z}$ is $s \times w^*-$closed in $F \times F^*$, for every $z \in A$ provided that T is $s \times w^*-$closed in $X \times X^*$.*

Proof. We may assume without loss of generality that $z = 0 \in A = F$. In this case $T_{A,z} = T_F$. Let $L : (X \times X^*, s \times w^*) \to (X \times F^*, s \times w^*)$, $L(x, x^*) = (x, x^*|_F)$. Then L is linear bounded and $\operatorname{Graph} T = L^{-1}(\operatorname{Graph} T_F)$ since $T = T + N_F$. Therefore T is $s \times w^*-$closed in $X \times X^*$ whenever T_F is $s \times w^*-$closed in $F \times F^*$.

Assume that F is closed and T is $s \times w^*-$closed in $X \times X^*$. Let $\Pi : (X \times X^*, s \times w^*) \to (X \times X^*, s \times w^*)/\{0\} \times F^\perp$, $\Pi(x, x^*) = (x, x^*) + \{0\} \times F^\perp$, be the projection map of $(X \times X^*, s \times w^*)$ onto the quotient space $(X \times X^*, s \times w^*)/\{0\} \times F^\perp$. Note that $\Pi^{-1}(\Pi(\operatorname{Graph} T)) = \operatorname{Graph} T + \{0\} \times F^\perp = \operatorname{Graph} T$ is $s \times w^*-$closed, that is, $\Pi(\operatorname{Graph} T)$ is closed in $(X \times X^*, s \times w^*)/\{0\} \times F^\perp$. The locally convex spaces $(X \times X^*, s \times w^*)/\{0\} \times F^\perp$ and $(X \times F^*, s \times w^*)$

are isomorphic via $j((x,x^*)+\{0\}\times F^\perp)=(x,x^*|_F)$ since F is closed. Using $L=j\circ\Pi$ we get that $\operatorname{Graph} T_F=L(\operatorname{Graph} T)=j(\Pi(\operatorname{Graph} T))$ is $s\times w^*-$closed in $F\times F^*$. ■

Proposition 7 *Let $(X,\|\cdot\|)$ be a normed space, let $T:X\rightrightarrows X^*$, let $A\subset X$ be an affine set such that $D(T)\subset A$, and let F be the linear subspace parallel to A. If T is demiclosed in $X\times X^*$ then $T_{A,z}$ is demiclosed in $F\times F^*$, for every $z\in A$. If, in addition, $T=T+N_A$, then T is demiclosed in $X\times X^*$ whenever $T_{A,z}$ is demiclosed in $F\times F^*$, for some $z\in A$.*

Proof. Up to a translation we may assume without loss of generality that $z=0\in A=F$. In this case $T_{A,z}=T_F$.

Assume that T is demiclosed in $X\times X^*$. Let $\{(x_i,f_i^*)\}_{i\in I}\subset T_F$ be such that $(x_i,f_i^*)\to(x,f^*)$, $s\times w^*$ in $F\times F^*$ and $(f_i^*)_i$ is (strongly) bounded in F^*. According to the Hahn-Banach Theorem, for every $i\in I$ there is $x_i^*\in X^*$ such that $x_i^*|_F=f_i^*$ and $\|x_i^*\|=\|f_i^*\|$. Hence $(x_i^*)_i$ is bounded in X^* and, eventually on a subnet, denoted by the same index for simplicity, $(x_i,x_i^*)\to(x,x^*)$, $s\times w^*$ in $X\times X^*$. This yields that $x^*|_F=f^*$ and $x^*\in Tx$ because T is demiclosed; whence $f^*\in T_Fx$. Therefore T_F is demiclosed in $F\times F^*$.

Assume that $T=T+N_F$ and T_F is demiclosed in $F\times F^*$. Let $\{(x_i,x_i^*)\}_{i\in I}\subset T$ be such that $(x_i,x_i^*)\to(x,x^*)$, $s\times w^*$ in $X\times X^*$ and $(x_i^*)_i$ is bounded in X^*. Then $(x_i,x_i^*|_F)\to(x,x^*|_F)$, $s\times w^*$ in $F\times F^*$ and $(x_i^*|_F)_i$ is bounded in F^*. Due to the demiclosedness of T_F we find that $x^*|_F\in T_Fx$, that is, there is $y^*\in Tx$ such that $x^*|_F=y^*|_F$. This implies $x^*\in y^*+F^\perp\subset Tx$. ■

Remark 5 *Under all the assumptions in Proposition 7, T_F is demiclosed in F^* on a subset $S\subset\overline{D(T)}$ iff T is demiclosed in X^* on S (that is, if $\{(x_i,x_i^*)\}_i\subset T$, $(x_i,x_i^*)\to(x,x^*)$ $s\times w^*$ in $X\times X^*$ and $x\in S$ then $(x,x^*)\in T$).*

Corollary 8 *Let X be a normed space, let $T:X\rightrightarrows X^*$ be such that $\operatorname{aff} D(T)$ is closed and $T=T+N_{\operatorname{aff} D(T)}$, and let F be the linear subspace parallel to $\operatorname{aff} D(T)$. Then*

(i) *T has w^*-closed values in X^* iff $T_{\operatorname{aff} D(T),z}$ has w^*-closed values in F^*, for every (some) $z\in\operatorname{aff} D(T)$;*

(ii) *T is $s\times w^*-$closed in $X\times X^*$ iff $T_{\operatorname{aff} D(T),z}$ is $s\times w^*-$closed in $F\times F^*$, for every (some) $z\in\operatorname{aff} D(T)$;*

(iii) *T is demiclosed in $X\times X^*$ iff $T_{\operatorname{aff} D(T),z}$ is demiclosed in $F\times F^*$, for every (some) $z\in\operatorname{aff} D(T)$.*

In the case of a finite-dimensional affine set passing through z and being spanned by the linearly independent set of directions $\{v_1,v_2,\ldots,v_d\}$

$$A:=A(z;v_1,v_2,..,v_d):=\{x=z+t_1v_1+..+t_dv_d\mid(t_1,..,t_d)\in\mathbb{R}^d\},\tag{5}$$

we associate to $T_{A,z}$ the finite-dimensional operator $\mathscr{T}_{A,z}:\mathbb{R}^d\rightrightarrows\mathbb{R}^d$ given by

$$(s_1,..,s_d)\in\mathscr{T}_{A,z}(t_1,..,t_d)\text{ iff }\exists x^*\in T(z+t_1v_1+..+t_dv_d):\ s_i=\langle v_i,x^*\rangle,\ i=\overline{1,d}.\tag{6}$$

Note that $\mathscr{T}_{A,z} = \mathscr{I}^* T_{A,z} \mathscr{I}$, where $\mathscr{I} : \mathbb{R}^d \to F := \operatorname{span}\{v_1, .., v_d\}$ is the isomorphism given by $\mathscr{I}(t_1, .., t_d) := t_1 v_1 + .. + t_d v_d$. This latter operator identity provides

$$\varphi_{\mathscr{T}_{A,z}}((t_1, .., t_d), (s_1, .., s_d)) = \varphi_{T_{A,z}}(t_1 v_1 + .. + t_d v_d, x^*), \tag{7}$$

where $x^* \in F^*$ is uniquely determined by $s_i = \langle v_i, x^* \rangle$, $i = \overline{1, d}$.

Alternately, $\mathscr{T}_{A,z} = J^* T_z J : D(\mathscr{T}_{A,z}) := J^{-1}(D(T) - z) \subset \mathbb{R}^d \rightrightarrows \mathbb{R}^d$, where $J := \iota_F \circ \mathscr{J} : \mathbb{R}^d \to X$, $J(t_1, .., t_d) := t_1 v_1 + .. + t_d v_d$.

Notice also that $\mathscr{T}_{A,z_2} = (\mathscr{T}_{A,z_1})_{J^{-1}(z_2 - z_1)}$, for every $z_1, z_2 \in A$.

Lemma 9 *Let X be a normed space, $T : X \rightrightarrows X^*$, $z \in X$, $\{v_1, .., v_d\}$ be a linearly independent subset of X, $F = \operatorname{span}\{v_1, .., v_d\}$, and $A = A(z; v_1, .., v_d)$. Then*

(i) *$\mathscr{T}_{A,z}$ is NI iff $T_{A,z}$ is NI,*

(ii) *$\mathscr{T}_{A,z} \in \mathscr{M}(\mathbb{R}^d)$ iff $T_{A,z} \in \mathscr{M}(F)$ iff $T + N_A \in \mathscr{M}(X)$.*

In the sequel, for $v \neq 0$, $z \in X$, $L := L(z; v) := A(z; v)$ is the line passing through z with direction v, $T_{L,z}(tv) := \{x^*|_{\mathbb{R}v} \mid x^* \in T(z + tv)\}$, $t \in \mathbb{R}$, and $\mathscr{T}_{L,z} : \mathbb{R} \rightrightarrows \mathbb{R}$, $s \in \mathscr{T}_{L,z} t$ if there is $x^* \in T(z + tv)$ such that $\langle v, x^* \rangle = s$.

Similarly, the plane passing through z with linearly independent set of directions $\{v_1, v_2\}$ is given by $P := P(z; v_1, v_2) := A(z; v_1, v_2)$, $T_{P,z}(t_1 v_1 + t_2 v_2) = \{x^*|_{\operatorname{span}\{v_1, v_2\}} \mid x^* \in T(z + t_1 v_1 + t_2 v_2)\}$, $t_1, t_2 \in \mathbb{R}$, and $\mathscr{T}_{P,z} : \mathbb{R}^2 \rightrightarrows \mathbb{R}^2$, $(s_1, s_2) \in \mathscr{T}_{P,z}(t_1, t_2)$ if there is $x^* \in T(z + t_1 v_1 + t_2 v_2)$ such that $\langle v_1, x^* \rangle = s_1$, $\langle v_2, x^* \rangle = s_2$.

Proposition 10 *Let $(X, \|\cdot\|)$ be a Banach space, let $T \in \mathcal{M}(X)$ be such that $\operatorname{ri} D(T) \neq \emptyset$ and let $A \subset \operatorname{aff} D(T)$ be finite-dimensional affine with $A \cap \operatorname{ri} D(T) \neq \emptyset$.*

(i) *If T has w^*-closed values then $\mathscr{T}_{A,z}$ has closed values, for every $z \in A$.*

(ii) *If T is demiclosed then $\mathscr{T}_{A,z}$ is closed, for every $z \in A$.*

Proof. We may assume without loss of generality that $\operatorname{aff} D(T) = X$, otherwise we replace T by $T_{\operatorname{aff} D(T),z}$ and acknowledge Corollary 8 . This change does not affect $\mathscr{T}_{A,z}$.

Let $A = A(z; v_1, .., v_d)$ and $z_0 := z + t_1^0 v_1 + .. + t_d^0 v_d \in A \cap \operatorname{int} D(T)$. From the local boundedness of T at z_0 there are $M, r > 0$ such that $z_0 + rB_X \subset D(T)$ and $\|x^*\| \leq M$, for every $x^* \in T(z_0 + ru)$, $u \in B_X$.

(ii) Consider $\{(s_1^n, .., s_d^n), (t_1^n, .., t_d^n)\}_{n \geq 1} \subset \mathscr{T}_{A,z}$, that is, there exists $x_n^* \in T(z + t_1^n v_1 + .. + t_d^n v_d)$ such that $\langle v_i, x_n^* \rangle = s_i^n$, $i = \overline{1, d}$, $n \geq 1$. Assume that $\lim_{n \to \infty}((s_1^n, .., s_d^n), (t_1^n, .., t_d^n)) = ((s_1, .., s_d), (t_1, .., t_d))$.

The monotonicity of T provides

$$\langle (t_1^n - t_1^0) v_1 + .. + (t_d^n - t_d^0) v_d - ru, x_n^* - x^* \rangle \geq 0, \ \forall x^* \in T(z_0 + ru), u \in B_X, n \geq 1,$$

from which $r\|x_n^*\| \leq M(|t_1^n - t_1^0|\|v_1\| + .. + |t_d^n - t_d^0|\|v_d\| + r) + (t_1^n - t_1^0) s_1^n + .. + (t_d^n - t_d^0) s_d^n$, $n \geq 1$. Hence $\{x_n^*\}_{n \geq 1}$ is bounded. On a subnet denoted for simplicity by the same index, $x_n^* \to x^* \in T(z + t_1v_1 + .. + t_dv_d)$ weakly-star in X^*, since T is demiclosed. Let $n \to \infty$ in $\langle v_i, x_n^* \rangle = s_i^n$, $i = \overline{1,d}$, to get $\langle v_i, x^* \rangle = s_i$, $i = \overline{1,d}$, that is $(s_1, .., s_d) \in \mathscr{T}_{A,z}(t_1, .., t_d)$.

The argument of (i) proceeds similarly with $(t_1^n, .., t_d^n) = (t_1, .., t_d) \in D(\mathscr{T}_{A,z})$, $n \geq 1$. ■

Since every convex set in a finite dimensional space has a non-empty relative (algebraic) interior which is dense in the set, the following definition is a natural extension to a general topological vector space context for the nearly-convex notion. A set $S \subset X$ is called *nearly-convex* if there is a convex set C such that $\operatorname{ri} C \neq \emptyset$ and $C \subset S \subset \overline{C}$. Equivalently, S is nearly-convex iff $\operatorname{ri} S$ is non-empty convex and $S \subset \operatorname{cl}(\operatorname{ri} S)$. Indeed, directly, we know that $\operatorname{ri} C = \operatorname{ri} \overline{C}$ and $\operatorname{cl}(\operatorname{ri} C) = \overline{C}$ (see [11, Lemma 11A b), p. 59]) from which $\operatorname{ri} S = \operatorname{ri} C$ is non-empty convex and $S \subset \overline{C} = \operatorname{cl}(\operatorname{ri} S)$. Conversely, $C = \operatorname{ri} S$ fulfills all the required conditions.

Lemma 11 *Let X be a Banach space, let $T : X \rightrightarrows X^*$ be such that $D(T)$ is nearly-convex, and let $A = A(z; v_1, v_2, .., v_d) \subset \operatorname{aff} D(T)$ be such that $A \cap \operatorname{ri} D(T) \neq \emptyset$. Then $D(\mathscr{T}_{A,z})$ is nearly-convex and*

$$\operatorname{int} D(\mathscr{T}_{A,z}) = J^{-1}(\operatorname{ri} D(T) - z),\ \operatorname{cl} D(\mathscr{T}_{A,z}) = J^{-1}(\overline{D(T)} - z), \tag{8}$$

where $J(t_1, .., t_d) = t_1v_1 + .. + t_dv_d$, $(t_1, .., t_d) \in \mathbb{R}^d$.

Proof. We may assume without loss of generality that $\operatorname{aff} D(T) = X$ (otherwise we replace X by $F := \operatorname{aff} D(T) \ni 0$ and T by T_F; $\mathscr{T}_{A,z}$ being impervious to this change). Let $D := \operatorname{int} D(T) - z$ and $S := D(T) - z$. Then $J^{-1}(D)$ is non-empty open convex, $J^{-1}(S) = D(\mathscr{T}_{A,z})$, $\overline{D} = \overline{S}$, and $\operatorname{int} \overline{S} = D$ (see e.g. [11, Lemma 11A b), p. 59]).

Fix $\bar{t}$ with $J\bar{t} \in D$. Since $J^{-1}(D) \subset D(\mathscr{T}_{A,z}) \subset J^{-1}(\overline{D})$ and $J^{-1}(D) \subset \operatorname{int} J^{-1}(\overline{D})$, $\operatorname{cl} J^{-1}(D) \subset J^{-1}(\overline{D})$ (due to the continuity of J) in order to conclude it suffices to show that $\operatorname{int} J^{-1}(\overline{D}) \subset J^{-1}(D)$ and $J^{-1}(\overline{D}) \subset \operatorname{cl} J^{-1}(D)$.

Let $t \in J^{-1}(\overline{D})$. Since $\operatorname{int} \overline{D} = D$, $(1 - \lambda)\bar{t} + \lambda t \in J^{-1}(D)$, for $0 \leq \lambda < 1$. Let $\lambda \uparrow 1$ to get $t \in \operatorname{cl} J^{-1}(D)$. Hence $J^{-1}(\overline{D}) = \operatorname{cl} J^{-1}(D)$ followed by $\operatorname{int} J^{-1}(\overline{D}) = \operatorname{int}(\operatorname{cl} J^{-1}(D)) = J^{-1}(D)$. ■

Lemma 12 *Let X be a Banach space and let $T : X \rightrightarrows X^*$ be such that $D(T)$ is nearly-convex and $T = T + N_{D(T)}$. Then $\mathscr{T}_{A,z} = \mathscr{T}_{A,z} + N_{D(\mathscr{T}_{A,z})}$, for every finite-dimensional affine set $A \subset X$ with $A \cap \operatorname{ri} D(T) \neq \emptyset$, $z \in A \cap \operatorname{ri} D(T)$.*

Proof. Condition $T = T + N_{D(T)}$ provides $T_z = T_z + (N_{D(T)})_z = T_z + N_{D(T)-z}$, $z \in X$. Recall that $\mathscr{T}_{A,z} = J^*T_zJ$, where $A = A(z; v_1, .., v_d)$, $J : \mathbb{R}^d \to X$, $J\widehat{t} = t_1v_1 + .. + t_dv_d$, $\widehat{t} = (t_1, .., t_d)$. Hence

$$\mathscr{T}_{A,z}\widehat{t} = \mathscr{T}_{A,z}\widehat{t} + J^*N_{D(T)-z}J\widehat{t},\ \forall \widehat{t} \in D(\mathscr{T}_{A,z}) = J^{-1}(D(T) - z).$$

But, for every $\widehat{t} \in D(\mathscr{T}_{A,z})$ we have from (8) and by the chain rule (see e.g. [28, Theorem 2.8.3(vii), p. 123]), that

$$N_{D(\mathscr{T}_{A,z})}\widehat{t} = N_{\operatorname{cl} D(\mathscr{T}_{A,z})}\widehat{t} = N_{J^{-1}(\overline{D(T)}-z)}\widehat{t} = \partial(\iota_{\overline{D(T)}-z} \circ J)(\widehat{t}) = J^*N_{\overline{D(T)}-z}J\widehat{t},$$

since $0 \in {}^{ic}(A - \overline{D(T)})$. The proof is complete. ■

4 Finite-dimensional context characterizations

Recall the next sum rule for maximal monotone operators followed by two of its consequences.

Theorem 13 ([27, Corollary 4]) *Let X be a Banach space and $M, N \in \mathscr{M}(X)$. Assume that ${}^{ic}D(M), {}^{ic}D(N)$ are nonempty and $0 \in {}^{ic}(D(M) - D(N))$. Then $M + N \in \mathscr{M}(X)$.*

Proposition 14 *Let X be a Banach space. If $T \in \mathscr{M}(X)$, A is a finite-dimensional affine subset of X, and $A \cap {}^{ic}D(T) \neq \emptyset$ then $T + N_A \in \mathscr{M}(X)$.*

Proof. We may apply the previous theorem since ${}^{ic}A = A$, $\operatorname{aff}(D(T) - A) = \operatorname{aff} D(T) - A$ is closed since $\operatorname{aff} D(T)$ is closed and A is finite-dimensional, and $0 \in {}^{i}D(T) - A \subset {}^{i}(D(T) - A)$.

Indeed, for the latter inclusion let $x_0 \in {}^{ic}D(T)$, $a_0 \in A$. Every $x \in \operatorname{aff}(D(T) - A)$ has the form $x = u - a$, with $u \in \operatorname{aff} D(T)$, $a \in A$. Therefore there is $\rho > 0$ such that $\lambda u + (1-\lambda)x_0 \in D(T)$, for every $\lambda \in [0, \rho]$. Hence $\lambda x + (1-\lambda)(x_0 - a_0) = \lambda u + (1-\lambda)x_0 - (\lambda a + (1-\lambda)a_0) \in D(T) - A$, for every $\lambda \in [0, \rho]$, that is, $x_0 - a_0 \in {}^{ic}(D(T) - A)$. The proof is complete. ■

Proposition 15 *Let X be a Banach space. If $T \in \mathscr{M}(X)$, A is a closed affine subset of X, and $A \cap \operatorname{core} D(T) \neq \emptyset$ then $T + N_A \in \mathscr{M}(X)$.*

This paper is mainly concerned with the following converse of Proposition 14.

Given $T \in \mathcal{M}(X)$ with the property that $T + N_A \in \mathscr{M}(X)$ for every finite-dimensional affine A (especially lines and planes) such that $A \cap {}^{ic}D(T) \neq \emptyset$ what additional conditions on T are needed, such as the closedness of its graph or convexity of its values, in order to obtain the maximality of T?

Proposition 16 *Let X be a Banach space and let $T \in \mathcal{M}(X)$ be such that ${}^{ic}D(T) \neq \emptyset$. If $T + N_L \in \mathscr{M}(X)$ for every line $L \subset \operatorname{aff} D(T)$ such that $L \cap {}^{ic}D(T) \neq \emptyset$ then $T_{\operatorname{aff} D(T), z}$ is NI, for every $z \in \operatorname{aff} D(T)$ and $\operatorname{ri} D(T) = {}^{ic}D(T)$, $\overline{D(T)} = \operatorname{cl}({}^{ic}D(T))$ are convex sets. In particular, ${}^{ic}D(T)$, $D(T)$, $\overline{D(T)}$ have the same (convex) relative interior and closure.*

Proof. First assume that $\operatorname{aff} D(T) = X$. Let (x_0, x_0^*) be m.r. to T and let $v \in X$ be such that $L := L(x_0; v)$ cuts ${}^{ic}D(T) = \operatorname{core} D(T)$. Since $n^* \in N_L(x)$ iff $x \in L$ and $\langle v, n^* \rangle = 0$, it is easily checked that (x_0, x_0^*) is m.r to $T + N_L \in \mathscr{M}(X)$. Therefore, if (x_0, x_0^*) is m.r. to T then $x_0 \in D(T)$ and for every v such that $L(x_0; v) \cap \operatorname{core} D(T) \neq \emptyset$ there is $x^* \in Tx_0$ such that $\langle v, x_0^* \rangle = \langle v, x^* \rangle$.

In particular T is NI (see [25, (1)]) and in this case we know that $\mathcal{R} := [\psi_T = c]$ is the unique maximal monotone extension of T (see [26, Proposition 4 (iii)]).

Every $(x,x^*)\in\mathcal{R}$ is m.r. to T. Hence $x\in D(T)$ and for every w such that $L(x;w)\cap\operatorname{core}D(T)\neq\emptyset$ there is $y^*\in Tx$ such that $\langle w,x^*\rangle=\langle w,y^*\rangle$. Therefore $D(\mathcal{R})=D(T)$.

Since $\mathcal{R}\in\mathscr{M}(X)$ and $\operatorname{core}D(\mathcal{R})\neq\emptyset$ we have $\operatorname{int}D(T)=\operatorname{core}D(T)$ and $\operatorname{cl}D(T)=\operatorname{cl}D(\mathcal{R})=\operatorname{cl}(\operatorname{int}D(\mathcal{R}))=\operatorname{cl}(\operatorname{int}D(T))$ are convex sets (see [27, Corollary 3] or [18, Theorem 1]).

In the general case we may assume without loss of generality that $0\in\operatorname{aff}D(T)=:F$ otherwise we change T with T_z where $z\in\operatorname{aff}D(T)$. We use the first part of our proof for $T_F\in\mathcal{M}(F)$. To this end note first that $T_F+(N_L)_F=(T+N_L)_F$, where $(N_L)_F\in\mathscr{M}(F)$ is the normal cone to $L\subset F$. According to Lemma 1, $(T+N_L)_F\in\mathscr{M}(F)$, since $(T+N_L)+N_F=T+N_L\in\mathscr{M}(X)$, for every line $L\subset F$, such that $L\cap{}^{ic}D(T)\neq\emptyset$. Therefore $\operatorname{ri}D(T)={}^{ic}D(T)$, $\overline{D(T)}=\operatorname{cl}({}^{ic}D(T))$ are convex sets and $T_{F,z}=(T_F)_z=(T_z)_F$ is NI, for every $z\in F$. The sets ${}^{ic}D(T)$, $D(T)$, $\overline{D(T)}$ have the same relative interior due to $\operatorname{ri}\overline{D(T)}=\operatorname{ri}(\operatorname{cl}(\operatorname{ri}D(T)))=\operatorname{ri}D(T)$ (see e.g. [11, Lemma 11A b), p. 59]). ■

Theorem 17 *Let $T\in\mathcal{M}(\mathbb{R}^d)$ be such that* $\operatorname{core}D(T)\neq\emptyset$. *Then $T\in\mathscr{M}(X)$ iff*

(i) *$T+N_L\in\mathscr{M}(\mathbb{R}^d)$ for every line $L\subset\mathbb{R}^d$ such that $L\cap\operatorname{core}D(T)\neq\emptyset$,*

(ii) *T has closed convex values,*

(iii) *$Tx=Tx+N_{\overline{D(T)}}x$, for every $x\in D(T)\setminus\operatorname{core}D(T)$.*

Proof. The direct implication is clear since (i) follows from Proposition 14 and (ii), (iii) are usual properties of maximal monotone operators.

For the converse suppose that (i), (ii), (iii) hold. As seen in the proof of Proposition 16, if (x_0,x_0^*) is m.r. to T then $x_0\in D(T)$,

$$\forall v:\ L(x_0;v)\cap\operatorname{core}D(T)\neq\emptyset,\ \exists x^*\in Tx_0,\ \langle v,x_0^*\rangle=\langle v,x^*\rangle,\tag{9}$$

and $\operatorname{core}D(T)=\operatorname{int}D(T)=\operatorname{int}\overline{D(T)}$, $\overline{D(T)}$ are convex sets.

Assume that $x_0^*\notin Tx_0$. Since Tx_0 is closed convex, for some $y_0\in\mathbb{R}^d$

$$\langle y_0,x_0^*\rangle>\sigma_{Tx_0}(y_0):=\sup\{\langle y_0,x^*\rangle\mid\ x^*\in Tx_0\},\tag{10}$$

that is, $[\sigma_C<0]$ is non-empty, where $C:=Tx_0-x_0^*\not\ni 0$.

Relation (9) shows that $L(x_0;z)\cap\operatorname{int}D(T)=\emptyset$ for every $z\in[\sigma_C<0]$. Hence $x_0\notin\operatorname{int}D(T)$ and $z\notin\operatorname{int}T_{\overline{D(T)}}x_0=\cup_{h>0}\frac{1}{h}(\operatorname{int}D(T)-x_0)$ (see e.g. [1, Proposition 7, p. 169]), where $T_{\overline{D(T)}}x_0$ stands for the tangent cone to the closed convex set $\overline{D(T)}$ at x_0. Therefore $[\sigma_C<0]\cap\operatorname{int}T_{\overline{D(T)}}x_0=\emptyset$.

Note that $[\sigma_C<0]$ is a convex cone whose closure $[\sigma_C\leq 0]\subset T_{\overline{D(T)}}x_0$. Indeed, using $Tx_0=Tx_0+N_{\overline{D(T)}}x_0$, every $z\in[\sigma_C\leq 0]$ satisfies $\langle z,n^*\rangle\leq 0$, for every $n^*\in N_{\overline{D(T)}}x_0$, that is, $z\in(N_{\overline{D(T)}}x_0)^-=T_{\overline{D(T)}}x_0$ (see e.g. [1, Proposition 4, p. 168]).

Assume that for some $u_0^*,v^*\in\mathbb{R}^d$, $L(u_0^*;v^*):=\{u_0^*+tv^*\mid t\in\mathbb{R}\}\subset Tx_0$. Since T is monotone for every $u\in D(T)$, $u^*\in Tu$, $t\in\mathbb{R}$, we find $\langle x_0-u,u_0^*+tv^*-u^*\rangle_{\mathbb{R}^d}\geq 0$; whence $\langle x_0-u,v^*\rangle_{\mathbb{R}^d}=0$. Since $x_0-D(T)$

has a non-empty interior this yields that $v^* = 0$ and so $L(u_0^*; v^*)$ is a singleton. Therefore C does not contain lines.

According to [19, Corollary 13.4.2, p. 118], $\operatorname{int}(\operatorname{dom} \sigma_C) \neq \emptyset$ and so σ_C is continuous on $\operatorname{int}(\operatorname{dom} \sigma_C)$. Since $[\sigma_C < 0]$ is nonempty we find that $\operatorname{int}([\sigma_C < 0]) \neq \emptyset$. Indeed let $x \in [\sigma_C < 0]$ and $y \in \operatorname{int}(\operatorname{dom} \sigma_C)$. Then for some $0 \leq t < 1$, $x_t := tx + (1-t)y \in [\sigma_C < 0] \cap \operatorname{int}(\operatorname{dom} \sigma_C)$. Since σ_C is continuous at x_t it follows that $x_t \in \operatorname{int}([\sigma_C < 0])$. This yields the contradiction $[\sigma_C < 0] \cap \operatorname{int} T_{\overline{D(T)}} x_0 \neq \emptyset$. Therefore $x_0^* \in Tx_0$ and $T \in \mathscr{M}(\mathbb{R}^d)$. ■

Condition (iii) in the previous theorem is equivalent to $T = T + N_{\overline{D(T)}}$.

Theorem 18 *Let $T \in \mathcal{M}(\mathbb{R}^d)$ be such that ${}^{i}D(T) \neq \emptyset$. Then $T \in \mathscr{M}(\mathbb{R}^d)$ iff*
(i) *$T + N_L \in \mathscr{M}(\mathbb{R}^d)$ for every line $L \subset \operatorname{aff} D(T)$ such that $L \cap {}^{i}D(T) \neq \emptyset$,*
(ii) *T has closed convex values,*
(iii) *$T = T + N_{\overline{D(T)}}$.*

Proof. The direct implication is trivial while for the converse assume, without loss of generality, that $0 \in F := \operatorname{aff} D(T)$.

Conditions (i), (ii), (iii) are transmitted from T to T_F via the fact that for every line $L \subset F$, $T + N_L + N_F = T + N_L \in \mathscr{M}(\mathbb{R}^d)$ and so by Lemma 1, $(T + N_L)_F = T_F + N_L \in \mathscr{M}(F)$ (argument already seen in the proof of Proposition 16), (iii) is easily checked to be hereditary from T to T_F, and T_F has closed convex values because so has T, F is finite dimensional, and (iii) holds (see also Proposition 5). According to Theorem 17, $T_F \in \mathscr{M}(F)$.

Again (iii) and $N_F \subset N_{\overline{D(T)}}$ provide $T = T + N_F$. According to Lemma 2 $T \in \mathscr{M}(\mathbb{R}^d)$. ■

Theorem 17 allows us to re-demonstrate Löhne's characterization of maximal monotonicity for finite-dimensional operators (see [13, Theorem 3.4]).

Theorem 19 *An operator $T \in \mathscr{M}(\mathbb{R}^d)$ iff* (i) *$T \in \mathcal{M}(\mathbb{R}^d)$;* (ii) *$D(T)$ is nearly convex, that is, there is a convex set C such that $C \subset D(T) \subset \overline{C}$;* (iii) *$T$ has convex values;* (iv) *$T = T + N_{D(T)}$;* (v) *T is closed.*

More precisely, [13, Theorem 3.4 (iv)] states that $(Tx)_\infty = N_{D(T)}x$, for every $x \in D(T)$, where $(Tx)_\infty$ stands for the recession cone of Tx. Since $Tx + (Tx)_\infty = Tx$, it is straightforward that [13, Theorem 3.4 (iv)] implies our condition (iv). Therefore Theorem 19 has a slightly weaker assumption (iv) than [13, Theorem 3.4].

Theorem 19 is trivial for a singleton domain operator. The following characterization of maximal monotonicity for $1-$dimensional operators is a simplified version and is used in the proof of Theorem 19.

Theorem 20 *Let $U : \mathbb{R} \rightrightarrows \mathbb{R}$ be such that $\operatorname{int} D(U) =: (\alpha, \omega) \neq \emptyset$. Consider the assumption: (F) $U(\alpha) = U(\alpha) + \mathbb{R}_-$ whenever $\alpha \in D(U)$, $U(\omega) = U(\omega) + \mathbb{R}_+$ whenever $\omega \in D(U)$.*

TFAE:

(i) $U \in \mathscr{M}(\mathbb{R})$,

(ii) $U \in \mathcal{M}(\mathbb{R})$, *U has convex values, (F) holds, and U is closed,*

(iii) $U \in \mathcal{M}(\mathbb{R})$, *U has closed convex values, (F) holds, and*

$$\inf U((t, +\infty)) \le \sup U(t), \ \forall t \in [\alpha, \omega) \cap \mathbb{R}, \tag{11}$$

$$\sup U((-\infty, t)) \ge \inf U(t) \ \forall t \in (\alpha, \omega] \cap \mathbb{R}. \tag{12}$$

Here $U(A) := \cup_{x \in A} U(x)$, *for* $A \subset \mathbb{R}$. *In all these cases* $\overline{D(U)} = \mathrm{cl}(\mathrm{int}\, D(U)) = [\alpha, \omega] \cap \mathbb{R}$ *and (F) is equivalent to* $U = U + N_{D(U)}$.

Remark 6 *Note that condition* $U(\alpha) = U(\alpha) + \mathbb{R}_-$ *whenever* $\alpha \in D(U)$ *comes to* $U(\alpha) = (-\infty, \sup U(\alpha)]$ *while* $U(\omega) = U(\omega) + \mathbb{R}_+$ *whenever* $\omega \in D(U)$ *is equivalent to* $U(\omega) = [\inf U(\omega), +\infty)$. *Therefore in (ii) (respectively (iii)) it suffices for U to have (closed) convex values only on* $\mathrm{int}\, D(U)$.

For $U \in \mathcal{M}(\mathbb{R})$ *relations (11), (12) represent a simplification of the closedness condition for U and can be equivalently restated as equalities, namely*

$$\lim_{r \downarrow t} (\inf U(r)) = \inf_{r > t} (\inf U(r)) = \inf U((t, +\infty)) = \sup U(t), \ \forall t \in [\alpha, \omega) \cap \mathbb{R},$$

$$\lim_{\ell \uparrow t} (\sup U(\ell)) = \sup_{\ell < t} (\sup U(\ell)) = \sup U((-\infty, t)) = \inf U(t), \ \forall t \in (\alpha, \omega] \cap \mathbb{R},$$

since $t \to \inf U(t)$, $t \to \sup U(t)$ *are non-decreasing.*

More interestingly, (11) is equivalent to

$$\lim_{r \downarrow t} (\sup U(r)) = \inf_{r > t} (\sup U(r)) = \sup U(t), \ \forall t \in [\alpha, \omega) \cap \mathbb{R},$$

while

$$\lim_{\ell \uparrow t} (\inf U(\ell)) = \sup_{\ell < t} (\inf U(\ell)) = \inf U(t), \ \forall t \in (\alpha, \omega] \cap \mathbb{R},$$

is an equivalent reformulation of (12).

Also, notice that (11) for $t = \alpha$ *spells* $\inf R(U) = -\infty$ *when* $\alpha \notin D(U)$ *or* $\sup U(\alpha) = \inf U((\alpha, +\infty))$ *when* $\alpha \in D(U)$. *Similarly (12) for* $t = \omega$ *spells* $\sup R(U) = +\infty$ *when* $\omega \notin D(U)$ *or* $\inf U(\omega) = \sup U((-\infty, \omega))$ *when* $\omega \in D(U)$.

Proof. (i) ⇒ (ii) Condition $U = U + N_{D(U)}$ together with $N_{(\alpha,\omega)}(\alpha) = \mathbb{R}_-$, $N_{(\alpha,\omega)}(\omega) = \mathbb{R}_+$(whenever α, ω are finite) provide (F). The other properties in (ii) are usual for $U \in \mathscr{M}(\mathbb{R})$.

(ii) ⇒ (iii) It suffices to verify (11), (12). Note that Ux is bounded, for every $x \in \mathrm{int}\, D(U)$, whenever $U \in \mathcal{M}(\mathbb{R})$. Clearly, (11) holds when $\inf U((t, \infty)) = -\infty$. Otherwise, for every $t \in [\alpha, \omega) \cap \mathbb{R}$, $\inf U((t, \infty))$ is finite. Take $(r_n)_n \subset (\alpha, \omega)$, $r_n \downarrow t$, $s_n := \min U(r_n)$ (attained since $U(r_n)$ is a closed bounded interval) such that $\inf U((t, \infty)) = \lim_{n \to \infty} s_n =: s \in \mathbb{R}$. Since U is closed, $s \in U(t)$ and so $s \le \sup U(t)$, that is, (11) holds. Note parenthetically that (11) holds for $t = \omega \in D(U)$ since $U(\omega) = U(\omega) + \mathbb{R}_+$ and $\sup U(\omega) = +\infty$. Relation (12) is verified similarly.

(iii) $\Rightarrow$ (i) First we show that $\omega < \infty$ implies $\sup R(U) = +\infty$. Indeed, assume that ω is finite. If $\omega \in D(U)$ from $U(\omega) = U(\omega) + \mathbb{R}_+$ it is clear that $\sup R(U) = +\infty$. If $\omega \notin D(U)$ then, according to (12), $\sup U((-\infty, \omega)) = \sup R(U) \geq \inf U(\omega) = \inf \emptyset = +\infty$. Similarly $\alpha > -\infty$ implies $\inf R(U) = -\infty$.

Let (t, s) be m.r. to U. If $t > \omega$ or $t = \omega \notin D(U)$ then ω is finite, and by the previous argument we find the contradiction $s \geq \sup R(U) = +\infty$. Therefore $t < \omega$ or $t = \omega \in D(U)$. Similarly, $t < \alpha$ or $t = \alpha \notin D(U)$ is impossible which leads to $t > \alpha$ or $t = \alpha \in D(U)$. Hence $t \in D(U) \cap [\alpha, \omega]$. In particular, since $U \in \mathcal{M}(\mathbb{R})$, we also get that $\overline{D(U)} = [\alpha, \omega] \cap \mathbb{R}$.

If $t \in (\alpha, \omega)$ then, due to (11), (12), we get $\inf U(t) \leq \sup U((-\infty, t)) \leq s \leq \inf U((t, \infty)) \leq \sup U(t)$. This yields $s \in U(t)$, since $U(t)$ is a closed interval. If $t = \alpha \in D(U)$ then, according to (11), $s \leq \inf U((\alpha, \infty)) \leq \sup U(\alpha)$; whence $s \in U(\alpha)$, since $U(\alpha) = (-\infty, \sup U(\alpha)]$. Similarly, $t = \omega \in D(U)$ provides $s \in U(\omega)$. The proof is complete. $\blacksquare$

Proof of Theorem 19. If $T \in \mathscr{M}(\mathbb{R}^d)$ then (iii), (iv), (v) are usual properties mainly due to the preservation of monotonicity. Since every finite-dimensional convex set has a non-empty convex relative (algebraic) interior which is dense in the set (see e.g. [20, Proposition 2.40, p. 64]), for (ii) notice that $\overline{D(T)}$ is convex and $\emptyset \neq C := \mathrm{ri}\, \overline{D(T)} \subset D(T) \subset \overline{C} = \overline{D(T)}$ (see also [20, Theorems 12.37, 12.41]).

For the converse note first that (ii) provide ${}^iD(T) = {}^iC = \mathrm{ri}\, C = \mathrm{ri}\, D(T)$, $\overline{D(T)} = \overline{C}$ are non-empty convex and T has closed convex values due to (iii) and (v).

Assume first that $\mathrm{aff}\, D(T) = \mathbb{R}^d$. Hence ${}^iD(T) = \mathrm{int}\, D(T)$. To conclude it suffices to check condition (i) in Theorem 17, that is, $T + N_L \in \mathscr{M}(\mathbb{R}^d)$ for every line $L \subset \mathbb{R}^d$ such that $L \cap \mathrm{int}\, D(T) \neq \emptyset$. According to Lemma 9 (ii), we need to show that $\mathscr{T}_{L,z} \in \mathscr{M}(\mathbb{R})$ for some $z \in \mathrm{int}\, D(T) \cap L$. To this end we prove that $U := \mathscr{T}_{L,z}$ verifies the conditions of Theorem 20 (ii).

Recall that $U = J^* T_z J$, where $L = L(z, v)$, $Jt = tv$, $t \in \mathbb{R}$ and $D(U) = J^{-1}((D(T) - z) \cap \mathbb{R}v)$. Therefore U has convex values since so does T. Because $z \in \mathrm{int}\, C$ we know that $\overline{(C - z) \cap \mathbb{R}v} = (\overline{C} - z) \cap \mathbb{R}v$. This yields that $(D(T) - z) \cap \mathbb{R}v$ is nearly convex and connected. Since $J : \mathbb{R} \to \mathbb{R}v$ is an isomorphism, $D(U)$ is a non-degenerate interval (and it is not a singleton because it contains the non-empty open set $J^{-1}(\mathrm{int}\, C - z)$).

Let $(\alpha, \omega) := \mathrm{int}\, D(U)$. Then $\alpha < 0$, $\omega > 0$ and, whenever α, ω are finite, $z + \alpha v, z + \omega v \in \overline{D(T)} \setminus \mathrm{int}\, D(T)$. Assume that $\alpha \in D(U)$, i.e., $z + \alpha v \in D(T)$. Take $n^* \in N_{D(T)}(z + \alpha v)$ such that $\langle z + \alpha v - y, n^* \rangle > 0$, for every $y \in \mathrm{int}\, D(T)$. Pick $y = z + \frac{\alpha}{2} v$ to get $\langle v, n^* \rangle < 0$. For every $s \in U(\alpha)$ there is $x^* \in T(z + \alpha v)$ such that $s = \langle v, x^* \rangle$. For every $\lambda < 0$ let $t = \lambda / \langle v, n^* \rangle > 0$. Then $x^* + tn^* \in T(z + \alpha v)$ due to (iv). Therefore $s + \lambda \in U(\alpha)$ and so $U(\alpha) = U(\alpha) + \mathbb{R}_-$. The second part of (F) is proved similarly.

Consider $s_n \in Ut_n$, i.e., $s_n = \langle v, x_n^* \rangle$ for some $x_n^* \in T(z + t_n v)$, $n \geq 1$, with $\lim_{n\to\infty}(s_n, t_n) = (s, t)$. From the local boundedness of T at z (see e.g. [2, Theorem 2]), there are $M, r > 0$ such that $z + rB \subset D(T)$ and $\|x^*\| \leq M$ for every $x^* \in T(z + ru)$, $u \in B := \{x \in \mathbb{R}^d \mid \|x\| \leq 1\}$. The monotonicity of T provides $\langle t_n v - ru, x_n^* - x^* \rangle \geq 0$, for every $x^* \in T(z + ru)$, $u \in B$, $n \geq 1$. This yields $r\|x_n^*\| \leq M(|t_n|\|v\| + r) + t_n s_n$, $n \geq 1$; whence $(x_n^*)_n$ is bounded. On a subsequence, denoted for simplicity by the same index, $x_n^* \to x_0^*$ in $\mathbb{R}^d$ and $x_0^* \in T(z + tv)$ since T is closed. Passing to limit in $s_n = \langle v, x_n^* \rangle$ we find $s = \langle v, x_0^* \rangle$, that is, $s \in Ut$ and so U is closed.

According to Theorem 20 (ii), $\mathscr{T}_{L,z} \in \mathscr{M}(\mathbb{R})$ and by the above argument $T \in \mathscr{M}(\mathbb{R}^d)$.

In the general case we may assume that $0 \in F := \operatorname{aff} D(T)$. Then T_F fulfills trivially (i), (ii), (iii), (iv) is due to $(N_{D(T)})_F$ being the normal cone to $D(T) \subset F$, and (v) follows from Corollary 8 (ii). Thus $T_F \in \mathscr{M}(F)$ followed by $T \in \mathscr{M}(\mathbb{R}^d)$ (recall that (iv) yields $T = T + N_F$). ■

5 Line-plane characterizations

Since the previous section provided several characterizations for the maximal monotonicity of operators defined in $\mathbb{R}$, the Banach spaces considered in this section are assumed to have dimension greater than one.

Proposition 21 *Let X be a Banach space and let $T \in \mathcal{M}(X)$ be such that ${}^{ic}D(T) \neq \emptyset$. If $T + N_P \in \mathscr{M}(X)$ for every plane P with $P \cap {}^{ic}D(T) \neq \emptyset$ then $T + N_L \in \mathscr{M}(X)$ for every line L with $L \cap {}^{ic}D(T) \neq \emptyset$.*

Proof. Let L be a line with with $L \cap {}^{ic}D(T) \neq \emptyset$ and P be a plane with $L \subset P$. Since $L \cap {}^{ic}(D(T) \cap P) \neq \emptyset$, by Proposition 14, we find that $T + N_L = (T + N_P) + N_L \in \mathscr{M}(X)$. ■

Theorem 22 *Let X be a Banach space and let $T \in \mathcal{M}(X)$ be such that ${}^{ic}D(T) \neq \emptyset$. Then $T \in \mathscr{M}(X)$ iff*

(i) *$T + N_P \in \mathscr{M}(X)$ for every plane $P \subset X$ such that $P \cap {}^{ic}D(T) \neq \emptyset$,*

(ii) *T has w^*-closed convex values.*

Due to Proposition 21 condition (i) in the previous theorem can be restated as $T + N_A$ is maximal monotone, for every affine set A generated by at most two linearly independent vectors and such that $A \cap {}^{ic}D(T) \neq \emptyset$. Note that this latter condition does not require a dimensional restriction on X.

Proof. For the direct implication (i) follows from Proposition 14 and (ii) is usual for $T \in \mathscr{M}(X)$.

For the converse let (x_0, x_0^*) be m.r. to T and let $A = A(x_0; v_1, v_2)$ be any affine set through x_0 generated by v_1, v_2 that cuts ${}^{ic}D(T)$. It is easily checked that (x_0, x_0^*) is m.r. to $T + N_A$, since $n^* \in N_A(x)$ iff $x \in A$ and $\langle v_1, n^* \rangle = \langle v_2, n^* \rangle = 0$. But $T + N_A \in \mathscr{M}(X)$.

Therefore, if (x_0, x_0^*) is m.r. to T then $x_0 \in D(T)$ and

$$\forall v_1, v_2 : A(x_0; v_1, v_2) \cap {}^{ic}D(T) \neq \emptyset, \exists x^* \in Tx_0 : \langle v_i, x_0^* \rangle = \langle v_i, x^* \rangle, i = 1, 2. \quad (13)$$

Assume that $x_0^* \notin Tx_0$. Since Tx_0^* is w^*−closed convex, by a separation argument there is $v_1 \in X$ such that

$$\langle v_1, x_0^* \rangle > \sup\{\langle v_1, y^* \rangle \mid \ y^* \in Tx_0\}. \quad (14)$$

Let v_2 be such that $A := A(x_0; v_1, v_2)$ cuts ${}^{ic}D(T)$ and so $T + N_A \in \mathscr{M}(X)$. By (13), this yields an $x^* \in Tx_0$ such that $\langle v_1, x_0^* \rangle = \langle v_1, x^* \rangle$ contrary to (14). This contradiction comes from the assumption that $x_0^* \notin Tx_0$. Hence $x_0^* \in Tx_0$ and $T \in \mathscr{M}(X)$. ■

Remark 7 *After looking at Theorems 18, 22 (see also Theorem 24 below) one wonders whether condition (iii) in Theorem 18 is necessary; in other words whether condition (i) in Theorem 22 could be relaxed by changing the planes with lines.*

Consider $C := \{(x, y) \in \mathbb{R}^2 \mid x \leq 0, y \leq 0\}$ *and* $T : D(T) := C \subset \mathbb{R}^2 \rightrightarrows \mathbb{R}^2$, $T(x, y) = N_C(x, y)$, *for every* $(x, y) \in C \setminus \{(0, 0)\}$ *and* $T(0, 0) = \mathbb{R}_+(1, 1)$. *Then* $T \notin \mathscr{M}(\mathbb{R}^2)$ *since* $T \subsetneq N_C$ *(more precisely* $T(0, 0) \subsetneq N_C(0, 0) = \{(x, y) \in \mathbb{R}^2 \mid x \geq 0, y \geq 0\} = \mathbb{R}_+(1, 0) + \mathbb{R}_+(0, 1)$*),* T *has closed convex values, and* $\operatorname{int} D(T) \neq \emptyset$.

Moreover, $T + N_L = N_C + N_L \in \mathscr{M}(\mathbb{R}^2)$ *(see e.g. Proposition 15), for every line* L *that cuts* $\operatorname{int} D(T)$*, that is, condition (i) in Theorem 18 is fulfilled. Indeed, this is straightforward if* L *does not pass through the origin. If* L *passes through the origin note that* L *cuts* $\operatorname{int} D(T)$ *iff* L *is non-vertical and has a positive-slope, that is,* $L := \mathbb{R}(1, m)$*, for some* $m > 0$*. Then* $N_L(0, 0) = L^\perp = \mathbb{R}(-m, 1)$ *and* $(T + N_L)(0, 0) = \mathbb{R}_+(1, 1) + \mathbb{R}(-m, 1) = \mathbb{R}_+(1, 0) + \mathbb{R}_+(0, 1) + \mathbb{R}(-m, 1) = (N_C + N_L)(0, 0)$*, since it is easily verifiable that* $\mathbb{R}_+(1, 0) \cup \mathbb{R}_+(0, 1) \subset \mathbb{R}_+(1, 1) + \mathbb{R}(-m, 1)$*. This suffices in order to have* $T + N_L = N_C + N_L$.



This example shows that condition (iii) in Theorem 18 is essential and, moreover, it cannot be relaxed to $T = T + N_{\operatorname{aff} D(T)}$*. However, if the lines in Theorem 18 (i) are upgraded to planes then (iii) can be relaxed to* $T = T + N_{\operatorname{aff} D(T)}$.

Theorem 23 *Let X be a Banach space and let $T \in \mathcal{M}(X)$ be such that ${}^{ic}D(T) \neq \emptyset$. Then $T \in \mathscr{M}(X)$ iff*
(i) *$T + N_P \in \mathscr{M}(X)$ for every plane $P \subset \operatorname{aff} D(T)$ such that $P \cap {}^{ic}D(T) \neq \emptyset$,*
(ii) *T has w^*-closed convex values,*
(iii) *$T = T + N_{\operatorname{aff} D(T)}$.*

Proof. The direct implication is straightforward. For the converse assume without loss of generality that $0 \in \operatorname{aff} D(T) =: F$. Apply the converse of Theorem 22 for $T_F : F \rightrightarrows F^*$. For every plane $P \subset F$ such that $P \cap {}^{ic}D(T) \neq \emptyset$, $T + N_P + N_F = T + N_P \in \mathscr{M}(X)$. From Lemma 1 we see that $(T + N_P)_F = T_F + (N_P)_F \in \mathscr{M}(F)$, where $(N_P)_F \in \mathscr{M}(F)$ is the normal cone to $P \subset F$. Hence $T_F \in \mathscr{M}(F)$ since, by Proposition 5, T_F has w^*-closed values in F^*. Together with (iii) this yields $T \in \mathscr{M}(X)$ (see Lemma 2). ■

The natural question whether the planes in condition (i) of Theorems 22, 23 can be replaced by lines is answered in the next result; thereby providing an extension to the infinite-dimensional context for Theorem 18.

Theorem 24 *Let X be a Banach space and let $T \in \mathcal{M}(X)$ be such that ${}^{ic}D(T) \neq \emptyset$. Then $T \in \mathscr{M}(X)$ iff*
(i) *$T + N_L \in \mathscr{M}(X)$ for every line $L \subset \operatorname{aff} D(T)$ such that $L \cap {}^{ic}D(T) \neq \emptyset$,*
(ii) *T has w^*-closed convex values,*
(iii) *$T = T + N_{D(T)}$.*

Proof. The direct implication is clear. For the converse note that from (iii), Lemma 2, and Proposition 5 we may assume without loss of generality that $\operatorname{aff} D(T) = X$ (otherwise we replace T by T_F where $0 \in F := \operatorname{aff} D(T)$); whence ${}^{ic}D(T) = \operatorname{core} D(T)$. From (i) via Proposition 16 we know that $\operatorname{core} D(T) = \operatorname{int} D(T)$, $\overline{D(T)} = \operatorname{cl}(\operatorname{int} D(T))$ are non-empty convex; in particular $D(T)$ is nearly-convex.

According to Theorem 22, it is enough to prove that $T + N_P \in \mathscr{M}(X)$ for every plane P with $P \cap \operatorname{int} D(T) \neq \emptyset$. From Lemma 9 (ii) we know that $T + N_P \in \mathscr{M}(X)$ iff $\mathscr{T}_{P,z} \in \mathscr{M}(\mathbb{R}^2)$ for some $z \in P \cap \operatorname{int} D(T)$.

For this choice of P and z we plan to use Theorem 17 for $\mathscr{T}_{P,z}$.

According to (ii) and Proposition 10 (i), $\mathscr{T}_{P,z}$ has closed convex values.

In this case recall (8), i.e.,

$$\operatorname{int} D(\mathscr{T}_{P,z}) = J^{-1}(\operatorname{int} D(T) - z),\ \operatorname{cl} D(\mathscr{T}_{P,z}) = J^{-1}(\overline{D(T)} - z), \tag{15}$$

where $P = A(z; v_1, v_2)$, $J : \mathbb{R}^2 \to X$, $J(t_1, t_2) := t_1 v_1 + t_2 v_2$, and $D(\mathscr{T}_{P,z}) = J^{-1}(D(T) - z)$.

We know from (iii) and Lemma 12 that $\mathscr{T}_{P,z}\hat{t} = \mathscr{T}_{P,z}\hat{t} + N_{D(\mathscr{T}_{P,z})}\hat{t}$, for every $\hat{t} = (t_1, t_2) \in D(\mathscr{T}_{P,z})$. According to Theorem 17, $\mathscr{T}_{P,z} \in \mathscr{M}(\mathbb{R}^2)$ as soon as $\mathscr{T}_{P,z} + N_\ell \in \mathscr{M}(\mathbb{R}^2)$ for every line $\ell \subset \mathbb{R}^2$ such that $\ell \cap \operatorname{int} D(\mathscr{T}_{P,z}) \neq \emptyset$.

Let $\ell \subset \mathbb{R}^2$ be a line such that $\ell \cap \operatorname{int} D(\mathscr{T}_{P,z}) \neq \emptyset$. Consider the line $L := J(\ell) + z \subset P$ or equivalently $\ell = J^{-1}(L - z)$. Note that $\iota_\ell = \iota_{L-z} \circ J$. Since $\ell \cap \operatorname{int} D(\mathscr{T}_{P,z}) \neq \emptyset$ we know from (15) that L cuts $\operatorname{int} D(T)$. We may apply the chain rule [28, Theorem 2.8.3(viii), p. 123] to get $N_\ell = J^* N_{L-z} J$. This yields $\mathscr{T}_{P,z} + N_\ell = \mathscr{A}_{P,z} := J^* A_z J$, where $A := T + N_L \in \mathscr{M}(X)$ due to (i). Note that $A = A + N_P$. According to Lemma 9 (ii), $\mathscr{A}_{P,z} \in \mathscr{M}(\mathbb{R}^2)$. The proof is complete. $\blacksquare$

Theorems 18, 22, 23, 24 show that under interiority conditions the sum result contained in Theorem 13 ([27, Corollary 4]) cannot be further improved in the sense that T need be maximal monotone.

The advantage of a condition of type Theorem 24 (i) over the (demi)closedness of the graph of an operator is that this condition involving lines can be replaced by the hemiclosedness or the so called "closedness on line" condition, as seen in Theorem 20.

Let $C \subset X$ be a convex set with $\operatorname{ri} D(T) \neq \emptyset$. Denote by $T_C(x)$ the tangent cone to C at $x \in X$ and by $S_C(x) := \bigcup_{h>0} h(C - x)$ the cone spanned by $C - x$. Then $\operatorname{ri} T_C(x) = S_{\operatorname{ri} C}(x)$, for very $x \in \overline{C}$ (see [1, Proposition 7, p. 169]).

We are ready for the following generalization of [4, Theorem 1.2], [8, Lemma 2.2], [21, Theorem 40.2, p. 155] (see also [21, Remark 40.3]), and [24, Lemma 4.2].

Theorem 25 *Let X be a Banach space and let $T \in \mathcal{M}(X)$ be such that $D(T)$ is nearly-convex.*

Then $T \in \mathscr{M}(X)$ iff $T = T + N_{D(T)}$ and (one of) the following assumptions hold(s):

(H) *T has w^*-closed convex values and for every $x \in \overline{D(T)}$, $v \in \operatorname{ri} T_{\overline{D(T)}}(x)$*

$$\inf_{h>0} \inf_{x^* \in T(x+hv)} \langle v, x^* \rangle \leq \sup_{x^* \in T(x)} \langle v, x^* \rangle. \tag{16}$$

(D) *T has convex values and is demiclosed,*

(R) *T is representable.*

Remark 8 *Whenever $T \in \mathcal{M}(X)$, the second part of condition* (H) *in the previous theorem is equivalent to the stronger forms: for $x \in \overline{D(T)}, v \in S_{\operatorname{ri} D(T)}(x)$*

$$\lim_{h \downarrow 0} \inf_{x^* \in T(x+hv)} \langle v, x^* \rangle = \inf_{h>0} \inf_{x^* \in T(x+hv)} \langle v, x^* \rangle = \sup_{x^* \in T(x)} \langle v, x^* \rangle,$$

$$\lim_{h \downarrow 0} \sup_{x^* \in T(x+hv)} \langle v, x^* \rangle = \inf_{h>0} \sup_{x^* \in T(x+hv)} \langle v, x^* \rangle = \sup_{x^* \in T(x)} \langle v, x^* \rangle,$$

since $h \to \inf_{x^* \in T(x+hv)} \langle v, x^* \rangle$, $h \to \sup_{x^* \in T(x+hv)} \langle v, x^* \rangle$ are non-decreasing. The last equivalent condition holds a striking resemblance to [13, Lemma 2.2] (one of the key results used in the proof of [13, Theorem 3.4]).

Proof. Let $T \in \mathcal{M}(X)$ be such that $D(T)$ is nearly-convex. For every $x \in \overline{D(T)}$, $v \in S_{\operatorname{ri} D(T)}(x)$ the line $L = L(x; v) \subset \operatorname{aff} D(T)$ cuts $\operatorname{ri} D(T)$ at some $z = x + t_0 v$ with $t_0 > 0$. Let $U := \mathscr{T}_{L,x}$. Then $D(U)$ is a non-degenerate interval with 0 inside its interior when $x \in \operatorname{ri} D(T)$ and with 0 as its left endpoint when $x \in \overline{D(T)} \setminus \operatorname{ri} D(T)$ (see Lemma 11). Recall that $s \in U(t)$ iff $s = \langle v, x^* \rangle$, for some $x^* \in T(x + tv)$; whence $\inf U(t) = \inf_{x^* \in T(x+tv)} \langle v, x^* \rangle$, $\sup U(t) = \sup_{x^* \in T(x+tv)} \langle v, x^* \rangle$, and $\sup U(0) = \sup_{x^* \in T(x)} \langle v, x^* \rangle$.

It is clear that for $T \in \mathscr{M}(X)$ conditions $T = T + N_{D(T)}$, (D), and (R) are fulfilled. According to Proposition 14, $T + N_L \in \mathscr{M}(X)$ and from Lemma 9 (ii), $\mathscr{T}_{L,x} \in \mathscr{M}(\mathbb{R})$. Condition (H) follows from relation (11) in Theorem 20 (iii) for $t = 0$.

Conversely, note that (R) $\Rightarrow$ (D) and T has w^*-closed convex values whenever (H) or (D) holds. According to Theorem 24, it suffices to show that $T + N_L \in \mathscr{M}(X)$ for ever line $L \subset \operatorname{aff} D(T)$ with $L \cap \operatorname{ri} D(T) \neq \emptyset$. To fix notation, let $L = L(z; v)$ with $z \in \operatorname{ri} D(T)$. According to Lemma 9 (ii) we need to show that $U := \mathscr{T}_{L,z} \in \mathscr{M}(\mathbb{R})$. To this end we use Theorem 20. Both (H), (D) imply via Proposition 10 (i) that U has closed convex values. Also, $\operatorname{int} D(U) =: (\alpha, \omega) \ni 0$, because $z \in \operatorname{ri} D(T)$. Condition $T = T + N_{D(T)}$ transfers to U via Lemma 12 and it clearly yields that $U(\alpha) = U(\alpha) + \mathbb{R}_-$, whenever $\alpha \in D(U)$ and $U(\omega) = U(\omega) + \mathbb{R}_+$, whenever $\omega \in D(U)$, since $N_{[\alpha,\omega]}(\alpha) = \mathbb{R}_-$, $N_{[\alpha,\omega]}(\omega) = \mathbb{R}_+$, that is, (F) holds.

If (D) is true then, according to Proposition 10 (ii), U is closed. In this case the conditions in Theorem 20 (ii) are met so $U \in \mathscr{M}(\mathbb{R})$ and we are done.

If (H) holds, to conclude by using Theorem 20 (iii), it remains to prove that (11), (12) hold.

Note that $v \in S_{\operatorname{ri} D(T)}(z + tv)$ for $\alpha \leq t < \omega$ and $-v \in S_{\operatorname{ri} D(T)}(z + tv)$ for $\alpha < t \leq \omega$; while $z + tv \in \overline{D(T)}$, for every $t \in [\alpha, \omega] \cap \mathbb{R}$. According to (16), for every $\alpha \leq t < \omega$

$$\inf U((t, +\infty)) = \inf_{h>0} \inf_{x^* \in T(z+tv+hv)} \langle v, x^* \rangle \leq \sup_{x^* \in T(z+tv)} \langle v, x^* \rangle = \sup U(t).$$

and for every $\alpha < t \leq \omega$

$$\begin{aligned} \sup U((-\infty, t)) &= \sup_{\ell<t} \sup_{x^* \in T(z+\ell v)} \langle v, x^* \rangle \\ &= -\inf_{h>0} \inf_{x^* \in T(z+tv+h(-v))} \langle -v, x^* \rangle \geq -\sup_{x^* \in T(z+tv)} \langle -v, x^* \rangle = \inf U(t). \end{aligned}$$

■

A stronger form of condition (H) in the previous theorem has been proven in [21, Lemma 40.1(d)] for $T \in \mathscr{M}(X)$ on $\operatorname{int} D(T)$. Similar continuity properties of this form are studied in our sixth section.

If X is finite-dimensional then (D) together with the monotonicity of T provide (H) as seen in the proof of Theorem 19. Hence all versions of Theorem 25 are extensions of Theorem 19. It is clear that at least for $x \in \operatorname{int} D(T)$ the demiclosedness of T implies the second part in (H) due to the local boundedness of T at x.

Our next aim is to make conditions (H) and (D) in Theorem 25 as disjoint as possible.

Theorem 26 *Let X be a Banach space and let $T \in \mathcal{M}(X)$ be such that $D(T)$ is nearly-convex. Then $T \in \mathscr{M}(X)$ iff*
(C) $T = T + N_{D(T)}$,
(H') *T has w^*-closed convex values and for every $x \in \overline{D(T)} \setminus \operatorname{ri} D(T)$, $v \in \operatorname{ri} T_{\overline{D(T)}}(x)$*

$$\inf_{h>0} \inf_{x^* \in T(x+hv)} \langle v, x^* \rangle \le \sup_{x^* \in T(x)} \langle v, x^* \rangle, \tag{17}$$

(D') *T is demiclosed on* $\operatorname{ri} D(T)$.

Proof. Assume first that $\operatorname{aff} D(T) = X$. We plan to use Theorem 25, therefore it suffices to show that (16) holds for $x \in \operatorname{int} D(T)$ and for every $v \in S_{\operatorname{int} D(T)}(x) = X$. Assume by contradiction that for a fixed $x \in \operatorname{int} D(T)$ there is $v \in X$ such that (16) does not hold. Taking Remark 8 into consideration, this entails the existence of $\epsilon_0 > 0$ such that

$$\langle v, x^* \rangle \ge \langle v, y^* \rangle + \epsilon_0, \ \forall h > 0, \ x^* \in T(x+hv), \ y^* \in Tx. \tag{18}$$

Take $h_i \downarrow 0$ such that for every i, $x + h_i v$ belongs to the neighborhood of x on which T is bounded. Hence any net $(x_i^*)_i$ with $x_i^* \in T(x + h_i v)$ is bounded. Eventually on a subnet, denoted by the same index for simplicity, $x_i^* \to z^* \in Tx$ weakly-star in X^*, by the demiclosedness of T. We obtain a contradiction if we use (18) for $h = h_i$, $x^* = x_i^*$, and $y^* = z^*$ and pass to limit.

In general we may assume that $0 \in \operatorname{aff} D(T) =: F$. The above argument applies to T_F via Propositions 5, 7, because (C) provides $T = T + N_F$. Therefore $T_F \in \mathscr{M}(F)$ and $T \in \mathscr{M}(X)$ via Lemma 2. ■

Proposition 27 *Let X be a Banach space and let $T \in \mathcal{M}(X)$ be such that $\operatorname{core} D(T) \neq \emptyset$. If T is demiclosed and Tx is unbounded for every $x \in D(T) \setminus \operatorname{core} D(T)$ then T is NI, $\operatorname{int} D(T) = \operatorname{core} D(T)$, $\overline{D(T)}$ are convex; in particular $D(T)$ is nearly-convex.*

Proof. Fix $y \in \operatorname{core} D(T)$ and $y^* \in Ty$. For every $x \in X$, set $S_x := \{s \in [0,1] \mid tx + (1-t)y \in D(T), \forall t \in [0,s]\}$, $s_x := \sup S_x$, and $z_x := s_x x + (1 - s_x)y$. Then $0 < s_x \le 1$ and $z_x \in \overline{D(T)} \setminus \operatorname{core} D(T)$.

Assume by contradiction that T is not NI that is $\varphi_T(x, x^*) < c(x, x^*)$ for some $(x, x^*) \in X \times X^*$. Then $x \notin D(T)$ since $D(T) \times X^* \subset [\varphi_T \ge c]$ (see e.g. [23, Proposition 3.2 (i)]). For every $s \in S_x$, let $t := s/s_x \in (0, 1]$. Then $tz_x + (1-t)y \in D(T)$. Again, using $D(T) \times X^* \subset [\varphi_T \ge c]$ one gets

$$\begin{aligned}
&t\varphi_T(z_x, x^*) + (1-t)c(y, y^*) = t\varphi_T(z_x, x^*) + (1-t)\varphi_T(y, y^*) \\
&\ge \varphi_T(t(z_x, x^*) + (1-t)(y, y^*)) = \varphi_T(tz_x + (1-t)y, tx^* + (1-t)y^*) \\
&\ge c(tz_x + (1-t)y, tx^* + (1-t)y^*) \\
&= tc(z_x, x^*) + (1-t)c(y, y^*) - t(1-t)c(z_x - y, x^* - y^*).
\end{aligned}$$

This yields $\varphi_T(z, x^*) \ge c(z_x, x^*) - (1-t)c(z_x - y, x^* - y^*)$. After we let $s \to s_x$, i.e., $t \to 1$, we get $\varphi_T(z_x, x^*) \ge c(z, x^*)$, from which $z_x \neq x$.

Let M be a maximal monotone extension of $T \cup \{(x, x^*)\}$. Then from $x \in D(M)$, $y \in \operatorname{core} D(M) = \operatorname{int} D(M)$, and $z_x \neq x$ we know that $z_x \in \operatorname{int} D(M)$. Therefore M and T are locally bounded at z_x. The demiclosedness of T implies that $z_x \in D(T)$ so $T(z_x)$ is non-empty bounded in contradiction to one of our assumptions. Therefore T is NI.

In this case we know that $\mathcal{R} := [\psi_T = c]$ is the unique maximal monotone extension of T (see [26, Proposition 4 (iii)]), $D(\mathcal{R}) \subset \mathrm{cl}(\operatorname{conv} D(T))$ since $\operatorname{dom} \psi_T \subset \mathrm{cl}_{s \times w^*} \operatorname{conv}(\operatorname{Graph} T)$, $\operatorname{int} D(\mathcal{R}) = \operatorname{core} D(\mathcal{R}) \neq \emptyset$, and $D(\mathcal{R})$ is nearly-convex (see [27, Corollary 3] or [18, Theorem 1]).

Take $x \in \operatorname{int} D(\mathcal{R}) \setminus \overline{D(T)}$. Then $z_x \neq x$ and $z_x \in \operatorname{int} D(\mathcal{R})$ since $\operatorname{int} D(\mathcal{R})$ is convex and contains y. Therefore the operators $\mathcal{R}$ and T are locally bounded at z_x and $z_x \in D(T)$ because T is demiclosed. Thus we reach at the contradiction $z_x \in D(T) \setminus \operatorname{core} D(T)$ and $T(z_x)$ is non-empty bounded.

Hence $\operatorname{int} D(\mathcal{R}) \subset \overline{D(T)} \subset \overline{D(\mathcal{R})}$; whence $\overline{D(T)} = \overline{D(\mathcal{R})}$ is convex and $D(\mathcal{R}) \subset \overline{D(T)}$ because $D(\mathcal{R})$ is nearly-convex. Again, for every $x \in \operatorname{int} D(\mathcal{R})$, $\mathcal{R}$ and T are locally bounded at x and $x \in D(T)$ due to the demiclosedness of T, that is, $\operatorname{int} D(\mathcal{R}) \subset D(T)$. This yields that $\operatorname{int} D(T) = \operatorname{int} D(\mathcal{R}) = \operatorname{core} D(T)$ is convex and $\overline{D(T)} = \overline{D(\mathcal{R})} = \mathrm{cl}(\operatorname{int} D(T))$. ∎

Remark 9 *The unboundedness of an operator* $T : X \rightrightarrows X^*$ *at every* $x \in D(T) \setminus \operatorname{core} D(T)$ *does not transmit to its restriction* $T_{\operatorname{aff} D(T), z}$ *even though* $\operatorname{ri} D(T) \neq \emptyset$. *Indeed, take* $F \subset X$ *a proper closed linear subspace and* $T = F \times F^{\perp}$. *Then* $Tx = F^{\perp}$ *is unbounded, for every* $x \in F$ *while* $T_F = F \times \{0\}$, *i.e.,* $T_F x = \{0\}$, *for* $x \in F$.

Lemma 28 *Let* $(X, \|\cdot\|)$ *be a Banach space, let* $T \in \mathcal{M}(X)$ *be such that* $\operatorname{core} D(T) \neq \emptyset$, *and let* $x \in D(T) \setminus \operatorname{core} D(T)$. *Consider the conditions*

(i) Tx *is unbounded;*

(ii) $\sup\{\langle x - y, x^* \rangle \mid x^* \in Tx\} = +\infty$, *for every (some)* $y \in \operatorname{core} D(T)$.

Then (ii) $\Rightarrow$ (i). *If, in addition,* $\operatorname{int} D(T) = \operatorname{core} D(T)$ *then* (i) $\Rightarrow$ (ii).

Proof. (ii) $\Rightarrow$ (i) Let $y \in \operatorname{core} D(T)$ be such that $\sup\{\langle x - y, x^* \rangle \mid x^* \in Tx\} = +\infty$. Since $x \neq y$ and $\langle x - y, x^* \rangle \leq \|x - y\| \|x^*\|$, this gives $\sup\{\|x^*\| \mid x^* \in Tx\} = +\infty$.

(i) $\Rightarrow$ (ii) For every $y \in \operatorname{int} D(T)$ let $M, r > 0$ be such that $y + rB_X \in D(T)$ and $\|y^*\| \leq M$, for every $y^* \in T(y + ru)$, $u \in B_X$. The monotonicity of T implies that $\langle x - y - ru, x^* - y^* \rangle \geq 0$, for every $x^* \in Tx$, $y^* \in T(y + ru)$, $u \in B_X$. This yields $\langle x - y, x^* \rangle \geq r\langle u, x^* \rangle - \|x - y - ru\| \|y^*\|$, for every $x^* \in Tx$, $y^* \in T(y + ru)$, $u \in B_X$; followed by $\langle x - y, x^* \rangle \geq r\langle u, x^* \rangle - (\|x - y\| + r)M$, for every $x^* \in Tx$, $u \in B_X$. Pass to supremum over $u \in B_X$ to get $\langle x - y, x^* \rangle \geq r\|x^*\| - (\|x - y\| + r)M$, for every $x^* \in Tx$; whence $\sup\{\langle x - y, x^* \rangle \mid x^* \in Tx\} = +\infty$. ∎

The advantage of condition (ii) in the previous lemma is that it transmits to $T_{\operatorname{aff} D(T), z}$; thereby allowing to relativize Proposition 27.

Proposition 29 *Let X be a Banach space and let $T \in \mathcal{M}(X)$ be such that ${}^{ic}D(T) \neq \emptyset$ and $T = T + N_{\operatorname{aff} D(T)}$. If T is demiclosed and for every $x \in D(T) \setminus {}^{ic}D(T)$ there is $y \in {}^{ic}D(T)$ such that $\sup\{\langle x - y, x^* \rangle \mid x^* \in Tx\} = +\infty$ then $T_{\operatorname{aff} D(T),z}$ is NI, for every $z \in \operatorname{aff} D(T)$ and $D(T)$ is nearly-convex.*

Proof. If $\operatorname{aff} D(T) = X$ the conclusion follows from Proposition 27 and Lemma 28. In general we may assume that $0 \in F := \operatorname{aff} D(T)$ otherwise we replace T by T_z with $z \in \operatorname{aff} D(T)$. Note that T_F is demiclosed due to $T = T + N_F$ and Proposition 7. Also $\sup\{\langle x - y, f^* \rangle \mid f^* \in T_F x\} = \sup\{\langle x - y, x^* \rangle \mid x^* \in Tx\} = +\infty$ which allows the conclusion for T_F. Therefore T_F is NI and $D(T)$ is nearly convex. ■

The following results are versions of Theorems 25, 26 mainly by relaxing condition (C).

Theorem 30 *Let X be a Banach space and $T : X \rightrightarrows X^*$ be such that ${}^{ic}D(T) \neq \emptyset$. Then $T \in \mathscr{M}(X)$ iff $T = T + N_{\operatorname{aff} D(T)}$, T is representable, and for every $x \in D(T) \setminus {}^{ic}D(T)$ there is $y \in {}^{ic}D(T)$ such that $\sup\{\langle x - y, x^* \rangle \mid x^* \in Tx\} = +\infty$. In this case ${}^{ic}D(T) = \operatorname{ri} D(T)$ and $D(T)$ is nearly-convex.*

Proof. For the direct implication if $T \in \mathscr{M}(X)$ then T is representable with ${}^{ic}D(T) = \operatorname{ri} D(T)$ and $D(T)$ nearly-convex (see [27, Corollary 3]) and every $x \in D(T) \setminus \operatorname{ri} D(T)$ is a support point of $D(T)$, i.e., there is $u^* \in N_{D(T)}x$, $u^* \neq 0$. This makes Tx unbounded because $Tx = Tx + N_{D(T)}x$ and so $\sup\{\langle x - y, x^* \rangle \mid x^* \in Tx\} = +\infty$, for every $y \in {}^{ic}D(T)$.

Proposition 29 may be used for the converse implications since every representable operator is monotone demiclosed to find that T_F is NI and $D(T)$ is nearly-convex, where we may assume without loss of generality that $0 \in F := \operatorname{aff} D(T)$.

Let $h \in \mathcal{R}_T$. Consider $g(x, f^*) = \inf\{h(x, x^*) \mid x^*|_F = f^*\}$, $(x, f^*) \in F \times F^*$. According to [24, Theorem 5.1], $g \in \mathcal{R}_{T_F}$ since $h, h^{\square} \le \psi_T$ (see [24, Remark 3.6] or [27, (5)]) which yields $\operatorname{Pr}_X(\operatorname{dom} h^*) \subset \operatorname{Pr}_X(\operatorname{dom} \psi_T) \subset \overline{D(T)} \subset F$ because $D(T)$ is nearly convex. Hence T_F is representable and from [22, Theorem 2.3] we know that $T \in \mathscr{M}(F)$. Again, Lemma 2 provides $T \in \mathscr{M}(X)$. ■

In particular, the previous result can be used to reprove Theorem 25 under the (R) assumption.

Corollary 31 *Let X be a Banach space and $T : X \rightrightarrows X^*$ be such that $D(T)$ is algebraically open (that is, $D(T) = {}^{ic}D(T)$). Then $T \in \mathscr{M}(X)$ iff $T = T + N_{\operatorname{aff} D(T)}$ and T is representable. In this case ${}^{ic}D(T) = \operatorname{ri} D(T)$ and $D(T)$ is nearly-convex.*

Remark 10 *One wonders whether, in the previous results, the contribution of $T|_{D(T) \setminus {}^{ic}D(T)}$ could be avoided. Let $T \in \mathscr{M}(X)$ with $\operatorname{int} D(T) \neq \emptyset$. In general, would the NI type and representability transmit from T to $T|_{\operatorname{int} D(T)}$? The answer is negative. For example, take $C \subsetneq X$ closed convex with $\operatorname{int} C \neq \emptyset$.*

Then $N_C \in \mathscr{M}(X)$ *(so it is NI and representable) while* $N_C|_{\operatorname{int} C} = \operatorname{int} C \times \{0\}$ *is neither NI (because it is not unique;* $X \times \{0\}$ *and* N_C *being two different strict maximal monotone extensions of* $N_C|_{\operatorname{int} C}$*) nor representable (since otherwise, according to Corollary 31,* $N_C|_{\operatorname{int} C} \in \mathscr{M}(X)$*).*

In the next result we avoid condition (C) completely.

Theorem 32 *Let* X *be a Banach space and let* $T : X \rightrightarrows X^*$ *be dual-representable. Consider the conditions*

(i) $D(T)$ *is nearly-convex,*

(ii) $^{ic}D(T)$ *is non-empty convex, and* $D(T) \subset \operatorname{cl}(^{ic}D(T))$*,*

(iii) $^{ic}\operatorname{Pr}_X(\operatorname{dom} h) \neq \emptyset$*, for some* $h \in \mathcal{D}_T$*,*

(iv) $T \in \mathscr{M}(X)$*.*

Then (i) $\Leftrightarrow$ (ii) $\Leftrightarrow$ (iii) $\Rightarrow$ (iv)*.*

Proof. (i) $\Rightarrow$ (ii) is straightforward.

(ii) $\Rightarrow$ (iii) Every $h \in \mathcal{D}_T$ has $h \leq \psi_T$; therefore $^{ic}D(T) \subset D(T) \subset \operatorname{Pr}_X(\operatorname{dom} h) \subset \operatorname{Pr}_X(\operatorname{dom} \psi_T) \subset \overline{D(T)} = \operatorname{cl}(^{ic}D(T))$ since $\overline{D(T)}$ is convex. This implies that $\operatorname{aff}(\operatorname{Pr}_X(\operatorname{dom} h)) = \operatorname{aff} D(T)$ and $^{ic}D(T) = ^{ic}\operatorname{Pr}_X(\operatorname{dom} h) \neq \emptyset$.

(iii) $\Rightarrow$ (iv) It suffices to prove that T is NI. To this end we show that $x \in D(T)$ whenever $z = (x, x^*)$ is m.r. to T.

Eventually by making a translation we may assume without loss of generality that $0 \in {}^{ic}\operatorname{Pr}_X(\operatorname{dom} h)$.

Let $f \in \Lambda(\mathbb{R}^2)$ be given by $f(t,s) = \inf\{h(y, y^*) \mid (y, y^*) \in C(t,s)\}$, where $C : \mathbb{R}^2 \rightrightarrows Z$, $(y, y^*) \in C(t,s)$ iff $y = tx$, $\langle x, y^* \rangle = s$. Note that C is linear and $s_{\mathbb{R}^2} \times (s_X \times w^*_{X^*})$–closed. Since $h \in \Gamma(Z)$ and $D := \operatorname{Im} C - \operatorname{dom} h = (\mathbb{R}x - \operatorname{Pr}_X(\operatorname{dom} h)) \times X^*$ has $\operatorname{aff} D = (\mathbb{R}x - \operatorname{aff}(\operatorname{Pr}_X(\operatorname{dom} h)) \times X^*$ closed, because $\operatorname{aff}(\operatorname{Pr}_X(\operatorname{dom} h)$ is closed and $\mathbb{R}x$ is finite dimensional, we get $0 \in {}^{ic}D$.

According to [28, Theorem 2.8.6 (v)], one gets

$$f^*(s,t) = \min\{h^*(y^*, y^{**}) \mid (s,t) \in C^*(y^*, y^{**})\}, \ (s,t) \in \mathbb{R}^2.$$

Since $(s,t) \in C^*(y^*, y^{**})$ iff $y^{**} = tx$, $\langle x, y^* \rangle = s$ one gets $f^{\square}(t,s) = f^*(s,t) = \min\{h^{\square}(tx, y^*) \mid \langle x, y^* \rangle = s\}$. Note that since $f \geq c$, we have $f^{\square\square} = \operatorname{cl} f \geq c$ and $f^{\square} \in \mathscr{D}_{[f^{\square} = c]}$. From [5, Theorem 3.1] we know that $[f^{\square} = c] \in \mathscr{M}(\mathbb{R}^2)$.

Note that from $h^{\square} \in \mathscr{R}_T$ one finds that for every $(t,s) \in [f^{\square} = c]$ there is $y^* \in T(tx)$ such that $\langle x, y^* \rangle = s$. Since $z = (x, x^*)$ is m.r. to T and implicitly to (tx, y^*) we get $(1-t)(\langle x, x^* \rangle - s) = \langle x - tx, x^* - y^* \rangle \geq 0$, that is $(1, \langle x, x^* \rangle)$ is m.r. to $[f^{\square} = c]$. Hence there is $y^* \in T(x)$ such that $\langle x, y^* \rangle = \langle x, x^* \rangle$; in particular $x \in \operatorname{dom} T$.

(iii) $\Rightarrow$ (iv) Whenever (iii) happens we know that $T \in \mathscr{M}(X)$. From [27, Corollary 3] we have that $^{ic}\operatorname{Pr}_X(\operatorname{dom} h) = {}^{ic}D(T) = \operatorname{ri} D(T)$ and; whence $D(T)$ is nearly-convex. ■

Remark 11 *Under the assumption* $0 \in {}^{ic}\operatorname{Pr}_X(\operatorname{dom} h)$ *for some* $h \in \mathcal{D}_T$ *the implication (iii)* $\Rightarrow$ *(iv) has been previously proved in [14, Theorem 3.1]. The advantage of our argument for (iii)* $\Rightarrow$ *(iv) is that, besides its brevity, it works for* X *merely a locally convex space under the modified assumption that* ${}^{ib}\operatorname{Pr}_X(\operatorname{dom} h) \neq \emptyset$, *for some* $h \in \mathcal{D}_T$, *where for* $S \subset X$, ${}^{ib}S = {}^{i}S$ *if the linear subspace parallel to* $\operatorname{aff} S$ *is barrelled,* ${}^{ib}S = \emptyset$ *otherwise.*

6 Continuity properties

Proposition 33 *Let* $(X, \|\cdot\|)$ *be a normed barrelled space and let* $A \subset X^*$ *be non-empty,* w^*-*closed, and convex. Then* $\operatorname{int}(\operatorname{dom} \sigma_A) \neq \emptyset$ *iff there exist* $y \in X$, $\beta \in \mathbb{R}$ *such that* $\langle y, x^* \rangle \geq \|x^*\| + \beta$, *for every* $x^* \in A$.

Proof. Assuming that $x_0 \in \operatorname{int}(\operatorname{dom} \sigma_A)$, σ_A is continuous at x_0; whence, for some $r > 0$, $f(u) := \sigma_A(x_0 + u) \leq \gamma < \infty$, for every $u \in rB_X$ or equivalently, $f \leq I_{rB_X} + \gamma$. Therefore $f^*(x^*) = I_A(x^*) - \langle x_0, x^* \rangle \geq (I_{rB_X} + \gamma)^*(x^*) = r\|x^*\| - \gamma$, for $x^* \in X^*$. This implies $\langle y, x^* \rangle \geq \|x^*\| + \beta$, for every $x^* \in A$, where $y = -(1/r)x_0$, $\beta = -\gamma/r$.

Conversely, assume that, for some $y \in X$, $\beta \in \mathbb{R}$, $\langle y, x^* \rangle \geq \|x^*\| + \beta$, for every $x^* \in A$. Then $h(x^*) := I_A(x^*) + \langle y, x^* \rangle \geq g(x^*) := \|x^*\| + \beta$, for every $x^* \in X^*$, followed by $h^*(u) = \sigma_A(-y + u) \leq g^*(u) = I_{B_X}(u) - \beta$, for every $u \in X$, that is, $\sigma_A(-y + u) \leq -\beta$, for every $u \in B_X$. This yields $-y \in \operatorname{int}(\operatorname{dom} \sigma_A)$. ■

Lemma 34 *Let* $(X, \|\cdot\|)$ *be a normed barrelled space and let* $A \subset X^*$ *be non-empty,* w^*-*closed, and convex with* $[\sigma_A < 0] \neq \emptyset$. *Then* $\operatorname{int}[\sigma_A < 0] \neq \emptyset$ *iff* $\operatorname{int}(\operatorname{dom} \sigma_A) \neq \emptyset$.

Proof. While the direct implication is straightforward for the converse let $y \in \operatorname{int}(\operatorname{dom} \sigma_A)$, $x \in [\sigma_A < 0]$, and $r, M > 0$ such that $\sigma_A(y + ru) \leq M < \infty$, for every $u \in B_X$ (since σ_A is continuous at y). Fix $1 > t > M/(M - \sigma_A(x))$ and let $x_t := tx + (1 - t)y$. We have

$$\sigma_A(x_t + (1-t)ru) = \sigma_A(tx + (1-t)(y + ru)) \leq t\sigma_A(x) + (1-t)M < 0, \ \forall u \in B_X,$$

which shows that $x_t \in \operatorname{int}([\sigma_A < 0])$. ■

Proposition 35 *Let* $(X, \|\cdot\|)$ *be a normed barrelled space, let* $T \in \mathcal{M}(X)$ *be such that* $\operatorname{int} D(T) \neq \emptyset$, *and let* $x \in X$, $x^* \in X^*$ *be such that* Tx *is* w^*-*closed convex and* $x^* \notin Tx$. *Then* $\operatorname{int}[\sigma_{Tx} < x^*] \neq \emptyset$. *If, in addition,* $Tx = Tx + N_{D(T)}x$ *and* $D(T)$ *is nearly-convex then* $\operatorname{int}[\sigma_{Tx} < x^*] \subset \operatorname{int} T_{D(T)}x = S_{\operatorname{int} D(T)}(x)$.

Proof. If $x \notin D(T)$ then $\sigma_{Tx} = -\infty$, $[\sigma_{Tx} < x^*] = X$, and the conclusion is trivial. If $x \in D(T)$ let $A := Tx - x^*$. A simple separation argument shows that $[\sigma_A < 0] = [\sigma_{Tx} < x^*] \neq \emptyset$. According to Lemma 34 and Proposition 33, it suffices to show that for some $y \in X$, $\beta \in \mathbb{R}$ one has $\langle y, y^* - x^* \rangle \geq \|y^* - x^*\| + \beta$, for every $y^* \in Tx$. Equivalently, we need to show that there exist $y \in X$, $\beta \in \mathbb{R}$ such that

$$\langle y, y^* \rangle \geq \|y^*\| + \beta, \ \forall y^* \in Tx. \tag{19}$$

Fix $z \in \operatorname{int} D(T)$ and let $M, r > 0$ be such that $z + rB_X \subset D(T)$ and $\|z^*\| \leq M$, for every $z^* \in T(z + ru)$, $u \in B_X$. The monotonicity of T provides

$$\langle z + ru - x, z^* - y^* \rangle \geq 0, \ \forall z^* \in T(z + ru), u \in B_X, y^* \in Tx.$$

This yields $\langle x - z, y^* \rangle \geq r\|y^*\| - M(\|z - x\| + r)$, that is, (19) holds with $y = (1/r)(x - z)$, $\beta = -M/r(\|z - x\| + r)$.

If, in addition, $Tx = Tx + N_{D(T)}x$ and $D(T)$ is nearly-convex then $[\sigma_{Tx} < x^*] \subset (N_{D(T)}x)^- = T_{\overline{D(T)}}x = \operatorname{cl}(\operatorname{int} T_{\overline{D(T)}}x) = \operatorname{cl} S_{\operatorname{int} D(T)}(x)$. Therefore $\emptyset \neq \operatorname{int}[\sigma_{Tx} < x^*] \subset \operatorname{int}(\operatorname{cl} S_{\operatorname{int} D(T)}(x)) = S_{\operatorname{int} D(T)}(x)$. ■

Remark 12 *The previous results allow us to present a different argument for the converse of Theorem 24.*

Indeed, recall from Proposition 16 that $\mathcal{R} := [\psi_T = c]$ *is the unique maximal monotone extension of* T *and* $D(T) = D(\mathcal{R})$ *is nearly-convex. We may assume without loss of generality that* $\operatorname{aff} D(T) = X$*. It suffices to show that* $T = \mathcal{R}$*. Assume by contradiction that there is* $(x, x^*) \in \mathcal{R}$ *such that* $x^* \notin Tx \neq \emptyset$*. According to Proposition 35, condition* $T = T + N_{D(T)}$ *provides* $\emptyset \neq \operatorname{int}[\sigma_{Tx} < x^*] \subset S_{\operatorname{int} D(T)}(x)$*. Take* $v \in \operatorname{int}[\sigma_{Tx} < x^*]$*. Since* $v \in S_{\operatorname{int} D(T)}(x)$ *we know that* $L(x, v) \cap \operatorname{int} D(T) \neq \emptyset$*, so, as seen in the proof of Proposition 16, there is* $y^* \in Tx$ *such that* $\langle v, x^* \rangle = \langle v, y^* \rangle$ *in contradiction with* $v \in [\sigma_{Tx} < x^*]$*. Therefore* $T = \mathcal{R} \in \mathscr{M}(X)$*.*

The following conjecture is stated in [3, p. 21]: - Every maximal monotone operator with a non-empty domain interior and defined in a Banach space is strongly× bounded weakly-star closed. A stronger form of this conjecture, namely the closedness property with respect to the strong×weak-star topology, is known to hold for the normal cone to a closed convex set with non-empty interior (see e.g. [7, Corollary on p. 58]). The next results give a positive answer to the mentioned conjecture in a relaxed context.

Theorem 36 *Let* $(X, \|\cdot\|)$ *be a normed barrelled space and* $T \in \mathcal{M}(X)$ *be such that* $\operatorname{int} D(T) \neq \emptyset$*. Let* $\{(x_i, x_i^*)\}_{i \in I} \subset T$ *be a net indexed on the directed set* $(I, \leq)$ *such that* $x_i \to x_0$*, strongly in* X*. Then there exist* $\gamma > 0$, $\beta \in \mathbb{R}$, $i_0 \in I$ *such that the following "a priori" estimate holds*

$$\langle x_0, x_i^* \rangle \geq \gamma \|x_i^*\| + \beta \ \forall i \geq i_0. \tag{20}$$

As a consequence we have the following two cases

(i) *If* $\limsup_{i \in I} \langle x_0, x_i^* \rangle < \infty$ *then, for some index* $i' \in I$, $\{x_i^*\}_{i \geq i'}$ *is bounded in* X^**. If, in addition*

(a) T *is demiclosed then* $x_0 \in D(T)$*;*

(b) T *is demiclosed and* $x_i^* \to x_0^*$*, weakly-star in* X^* *then* $(x_0, x_0^*) \in T$*.*

In particular, every monotone demiclosed operator with a non-empty domain interior is strongly×weakly-star closed.
(ii) *If* $\limsup_{i\in I}\langle x_0, x_i^*\rangle = \infty$ *then, at least on a subnet,* $\|x_i^*\|^{-1}x_i^* \to u^* \in N_{D(T)}x_0$ *weakly-star in* X^**, and* $\langle x_0, u^*\rangle \geq \gamma$.

Proof. See the published version. ■

Theorem 37 *Let* X *be a Banach space and let* $T \in \mathcal{M}(X)$ *be demiclosed with* $\operatorname{ri} D(T) \neq \emptyset$ *and* $T = T + N_{\operatorname{aff} D(T)}$. *Then* T *is* $s \times w^*-$*closed in* $X \times X^*$.

Proof. Assume without loss of generality that $0 \in F := \operatorname{aff} D(T)$. According to Proposition 7, $T_F \in \mathcal{M}(F)$ is demiclosed and $\operatorname{int} D(T_F) = \operatorname{ri} D(T) \neq \emptyset$. From Theorem 36, T_F is $s \times w^*-$closed in $F \times F^*$. Proposition 6 completes the proof. ■

Theorem 38 *Let* X *be a Banach space and let* $T \in \mathscr{M}(X)$ *be such that* ${}^{ic}D(T) \neq \emptyset$. *Then* T *is* $s \times w^*-$*closed in* $X \times X^*$.

Proof. Again we may assume that $0 \in F := \operatorname{aff} D(T)$. Since $T \in \mathscr{M}(X)$ we know that T is demiclosed, $\operatorname{ri} D(T) = {}^{ic}D(T) \neq \emptyset$, and $T = T + N_F$. The conclusion follows from the previous theorem. ■

It is common knowledge that a maximal monotone operator with a non-empty domain interior and defined in a Banach space is strongly×weakly-star upper semi-continuous on the interior of its domain (see e.g. [12, Theorem 6.7, p. 55], [9, Proposition B, p. 113], or [6, Theorem 2.5 (i), p. 155]). For completeness we include a proof of this result in a slightly relaxed context.

Theorem 39 *Let* X *be a normed barrelled space and let* $T \in \mathcal{M}(X)$. *If* T *is demiclosed then* T *is* $s \times w^*-$*upper semicontinuous at every* $x \in \operatorname{core} D(T)$.

Proof. Assume that T is not $s \times w^*-$upper semicontinuous at some $x \in \operatorname{core} D(T)$, that is, there is a w^*-open set $V \supset Tx$, $x_n \to x$, strongly in X, and $x_n^* \in Tx_n$ such that $x_n^* \notin V$, for every $n \geq 1$. For n large enough, $x_n \in U$, where U is the neighborhood of x on which $T(U)$ is bounded. Therefore $(x_n^*)_n$ is bounded and eventually on a subnet, denoted by the same index for simplicity, $x_n^* \to x^*$ weakly-star in X^*. Then $x^* \in Tx$ since T is demiclosed and $x^* \notin V$ because V is w^*-open. This spells the obvious contradiction $Tx \not\subset V$. ■

It is easily verifiable that the $s \times w^*-$upper semicontinuity of a maximal monotone operator T does not necessarily hold on the boundary of $D(T)$ even though the context is finite dimensional. For example take B the closed unit ball in $\mathbb{R}^2$ endowed with the usual euclidean inner product "$\langle\cdot,\cdot\rangle$" and norm "$\|\cdot\|$". Then N_B is not upper semicontinuous at any $x \in \mathbb{R}^2$ with $\|x\| = 1$. Indeed, $N_Bx = \mathbb{R}_+x$, for every $\|x\| = 1$ and for every $t \geq 0$, $\|y\| = 1$, $d := \operatorname{dist}(ty, \mathbb{R}_+x) = t\sqrt{1 - \langle x, y\rangle^2}$. Hence for $W := \frac{1}{2}B$, $V := N_Bx + W$ (a neighborhood of N_Bx), and every U a neighborhood of x we pick $y \in U$ with

$\|y\| = 1$, $y \neq x$ (so $\langle x, y\rangle \neq 1$), and $t = (1 - \langle x, y\rangle^2)^{-1/2}$. Then $ty \in N_B(U)$ and $ty \notin V$ due to $\operatorname{dist}(ty, \mathbb{R}_+ x) = 1$.



However, in a finite dimensional settings a different form of the $s \times w^*-$upper semicontinuity, namely, the (Q) (or Cesari) property holds for maximal monotone operators (see e.g. [13, Lemma 3.2], [12]). Our final aim is to extend the (Q) property to an infinite dimensional context.

Recall that $T : X \rightrightarrows X^*$ *has property (Q)* (or is *upper* $\mathscr{C}-$*semicontinuous*) *at* $x \in X$ (with respect to the $s \times w^*-$topology on $X \times X^*$) if for every net $\{x_i\}_{i\in I} \subset X$ such that $x_i \to x$, strongly in X we have

$$\bigcap_{i\in I} \operatorname{cl}_{w^*}(\operatorname{conv} \bigcup_{j\geq i} Tx_j) \subset Tx.$$

Clearly, property (Q) (as well as the $s \times w^*-$usc property) at x has substance only when $x \in \overline{D(T)}$ and, at least on a subnet, $\{x_i\}_i \subset D(T)$. The operator T *has property (Q)* if it has property (Q) at each $x \in X$.

Theorem 40 *Let X be a normed barrelled space and let $T \in \mathscr{M}(X)$ be such that* $\operatorname{core} D(T) \neq \emptyset$. *Then T has property (Q).*

Proof. Let $x \in X$, let $\{x_i\}_{i\in I}$ be such that $x_i \to x$, strongly in X, and let $x^* \in \bigcap_{i\in I} \operatorname{cl}_{w^*}(\operatorname{conv} \bigcup_{j\geq i} Tx_j)$.

For every $i \in I$ there are $k := k_i$, $I \ni j_1, .., j_k \geq i$, $x^*_{j_1} \in Tx_{j_1}, .., x^*_{j_k} \in Tx_{j_k}$, $\lambda_1, .., \lambda_k > 0$ such that $\sum_{\ell=1}^k \lambda_\ell = 1$ and $y^* := \sum_{\ell=1}^k \lambda_\ell x^*_{j_\ell}$ satisfies $|\langle x, y^* - x^*\rangle| \leq 1$. Pick an index $p \in \{1, .., k_i\}$ such that $\langle x, x^*_{j_p}\rangle \leq \langle x, y^*\rangle$ (such an index exists since $\langle x, y^*\rangle = \sum_{\ell=1}^k \lambda_\ell \langle x, x^*_{j_\ell}\rangle$) and denote j_p by $\varphi(i)$. In this way we generate a map $\varphi : I \to I$ such that $\varphi(i) \geq i$, for every $i \in I$ and a net $\{(x_{\varphi(i)}, x^*_{\varphi(i)})\}_{i\in I} \subset T$ such that $\{x_{\varphi(i)}\}_{i\in I}$ is a subnet of $\{x_i\}_{i\in I}$ and $\sup_{i\in I}\langle x, x^*_{\varphi(i)}\rangle \leq \langle x, x^*\rangle + 1 < \infty$. According to Theorem 36 (i), $x \in D(T)$, since T is demiclosed.

Assume, by contradiction, that $x^* \notin Tx$. Under our assumptions we may apply Proposition 35 to get $v \in \operatorname{int}[\sigma_{Tx} < x^*] \subset \operatorname{int} T_{D(T)}x$. Let $\epsilon_0 := (\langle v, x^*\rangle - \sigma_{Tx}(v))/2 > 0$. As previously seen, for every $i \in I$ there are $k := k_i$, $I \ni j_1, .., j_k \geq i$, $x^*_{j_1} \in Tx_{j_1}, .., x^*_{j_k} \in Tx_{j_k}$, $\lambda_1, .., \lambda_k > 0$ such that $\sum_{\ell=1}^k \lambda_\ell = 1$ and

$y^* := \sum_{\ell=1}^{k} \lambda_\ell x^*_{j_\ell}$ satisfies $|\langle v, y^* - x^*\rangle| \le \epsilon_0$. Pick an index $p \in \{1,..,k_i\}$ such that $\langle v, x^*_{j_p}\rangle \ge \langle v, y^*\rangle$ and denote j_p by $\varphi(i)$ to generate the map $\varphi : I \to I$ with the properties $\varphi(i) \ge i$, for every $i \in I$, $\{(x_{\varphi(i)}, x^*_{\varphi(i)})\}_{i\in I} \subset T$, $x_{\varphi(i)} \to x$, strongly in X, and

$$\inf_{i\in I} \langle v, x^*_{\varphi(i)}\rangle \ge \langle v, x^*\rangle - \epsilon_0. \tag{21}$$

According to Theorem 36, there exist $\gamma > 0$, $\beta \in \mathbb{R}$, $i_0 \in I$ such that

$$\langle x, x^*_{\varphi(i)}\rangle \ge \gamma \|x^*_{\varphi(i)}\| + \beta \ \forall i \ge i_0. \tag{22}$$

If $\limsup_{i\in I} \langle x, x^*_{\varphi(i)}\rangle < \infty$ then, from Theorem 36 (i), $\{x^*_{\varphi(i)}\}_{i\ge i'}$ is bounded so, at least on a subnet, $x^*_{\varphi(i)} \to x^*_0 \in Tx$, weakly-star in X^*. Passing to limit in (21) leads to the contradiction $\sigma_{Tx}(v) \ge \langle v, x^*_0\rangle \ge \langle v, x^*\rangle - \epsilon_0 = \sigma_{Tx}(v) + \epsilon_0$.

Therefore, at least on a subnet, denoted for simplicity by the same index, $\lim_{i\in I} \langle x, x^*_{\varphi(i)}\rangle = \lim_{i\in I} \|x^*_{\varphi(i)}\| = \infty$. From Theorem 36 (ii) we know that $\|x^*_{\varphi(i)}\|^{-1} x^*_{\varphi(i)} \to u^* \in N_{D(T)}x$, weakly-star in X^*, and $\langle x, u^*\rangle \ge \gamma$. Divide (21) by $\|x^*_{\varphi(i)}\|$ and pass to limit to obtain $\langle v, u^*\rangle = 0$ due to the fact that $v \in T_{D(T)}x = (N_{D(T)}x)^-$.

Take $\delta > 0$ such that $v + \delta x \in T_{D(T)}x$. If $\limsup_{i\in I} \langle v + \delta x, x^*_{\varphi(i)}\rangle > -\infty$, i.e., at least on a subnet, $\{\langle v + \delta x, x^*_{\varphi(i)}\rangle\}_i$ is bounded from below: $\langle v + \delta x, x^*_{\varphi(i)}\rangle \ge C > -\infty$, for every $i \in I$. Divide again by $\|x^*_{\varphi(i)}\|$ and pass to limit to find the contradiction $\delta\gamma \le \langle v + \delta x, u^*\rangle = 0$. Therefore $\lim_{i\in I} \langle v + \delta x, x^*_{\varphi(i)}\rangle = -\infty$ from which we get, taking (21) into account, that $\lim_{i\in I} \langle x, x^*_{\varphi(i)}\rangle = -\infty$, which contradicts (22). The proof is complete. $\blacksquare$

Theorem 41 *Let X be a Banach space and let $T \in \mathscr{M}(X)$ be such that ${}^{ic}D(T) \ne \emptyset$. Then T has property (Q).*

Proof. As usual assume that $0 \in F := \operatorname{aff} D(T)$. Since $T \in \mathscr{M}(X)$ we know that $T_F \in \mathscr{M}(F)$, $\operatorname{int} D(T_F) = \operatorname{ri} D(T) = {}^{ic}D(T) \ne \emptyset$, and $T = T + N_F$. According to Theorem 40, T_F has property (Q) with respect to the $s \times w^*$−topology on $F \times F^*$.

Let $x \in \overline{D(T)}$, let $\{x_i\}_{i\in I} \subset D(T)$ be such that $x_i \to x$, strongly in X, and let $x^* \in \bigcap_{i\in I} \operatorname{cl}_{w^*}(\operatorname{conv} \bigcup_{j\ge i} Tx_j)$. Then $x^*|_F \in \bigcap_{i\in I} \operatorname{cl}_{w^*}(\operatorname{conv} \bigcup_{j\ge i} T_F x_j)$ due to the continuity of $\iota^* : (X^*, w^*) \to (F^*, w^*)$, $\iota^*(x^*) = x^*|_F$. Hence $x^*|_F \in T_F x$, that is $x^* \in Tx + F^\perp = Tx$. $\blacksquare$

References

[1] Jean-Pierre Aubin and Ivar Ekeland. *Applied nonlinear analysis.* Pure and Applied Mathematics (New York). John Wiley & Sons Inc., New York, 1984. A Wiley-Interscience Publication.

[2] Jon Borwein and Simon Fitzpatrick. Local boundedness of monotone operators under minimal hypotheses. *Bull. Austral. Math. Soc.*, 39(3):439–441, 1989.

[3] Jonathan M. Borwein. Asplund decompositions of monotone operators. In *CSVAA 2004—control set-valued analysis and applications*, volume 17 of *ESAIM Proc.*, pages 19–25. EDP Sci., Les Ulis, 2007.

[4] Felix E. Browder. Multi-valued monotone nonlinear mappings and duality mappings in Banach spaces. *Trans. Amer. Math. Soc.*, 118:338–351, 1965.

[5] Regina Sandra Burachik and B. F. Svaiter. Maximal monotonicity, conjugation and the duality product. *Proc. Amer. Math. Soc.*, 131(8):2379–2383 (electronic), 2003.

[6] Ioana Cioranescu. *Geometry of Banach spaces, duality mappings and nonlinear problems*, volume 62 of *Mathematics and its Applications*. Kluwer Academic Publishers Group, Dordrecht, 1990.

[7] Frank H. Clarke. *Optimization and nonsmooth analysis*. Canadian Mathematical Society Series of Monographs and Advanced Texts. John Wiley & Sons Inc., New York, 1983. A Wiley-Interscience Publication.

[8] S. Fitzpatrick and R. R. Phelps. Bounded approximants to monotone operators on Banach spaces. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 9(5):573–595, 1992.

[9] S. P. Fitzpatrick. Continuity of nonlinear monotone operators. *Proc. Amer. Math. Soc.*, 62(1):111–116 (1977), 1976.

[10] Simon Fitzpatrick. Representing monotone operators by convex functions. In *Workshop/Miniconference on Functional Analysis and Optimization (Canberra, 1988)*, volume 20 of *Proc. Centre Math. Anal. Austral. Nat. Univ.*, pages 59–65. Austral. Nat. Univ., Canberra, 1988.

[11] Richard B. Holmes. *Geometric functional analysis and its applications*. Springer-Verlag, New York, 1975. Graduate Texts in Mathematics, No. 24.

[12] Shui Hung Hou. On property (Q) and other semicontinuity properties of multifunctions. *Pacific J. Math.*, 103(1):39–56, 1982.

[13] Andreas Löhne. A characterization of maximal monotone operators. *Set-Valued Anal.*, 16(5-6):693–700, 2008.

[14] M. Marques Alves and B. F. Svaiter. Maximal monotonicity, conjugation and the duality product in non-reflexive Banach spaces. *J. Convex Anal.*, 17(2):553–563, 2010.

[15] George J. Minty. On the maximal domain of a "monotone" function. *Michigan Math. J.*, 8:135–137, 1961.

[16] Robert R. Phelps. *Convex functions, monotone operators and differentiability*, volume 1364 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1989.

[17] Robert R. Phelps. *Convex functions, monotone operators and differentiability*, volume 1364 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, second edition, 1993.

[18] R. T. Rockafellar. Local boundedness of nonlinear, monotone operators. *Michigan Math. J.*, 16:397–407, 1969.

[19] R. Tyrrell Rockafellar. *Convex analysis*. Princeton Landmarks in Mathematics. Princeton University Press, Princeton, NJ, 1997. Reprint of the 1970 original, Princeton Paperbacks.

[20] R. Tyrrell Rockafellar and Roger J.-B. Wets. *Variational analysis*, volume 317 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1998.

[21] Stephen Simons. *Minimax and monotonicity*, volume 1693 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1998.

[22] M. D. Voisei. A maximality theorem for the sum of maximal monotone operators in non-reflexive Banach spaces. *Math. Sci. Res. J.*, 10(2):36–41, 2006.

[23] M. D. Voisei. Calculus rules for maximal monotone operators in general Banach spaces. *J. Convex Anal.*, 15(1):73–85, 2008.

[24] M. D. Voisei. The sum and chain rules for maximal monotone operators. *Set-Valued Anal.*, 16(4):461–476, 2008.

[25] M. D. Voisei. A sum theorem for (FPV) operators and normal cones. *J. Math. Anal. Appl.*, 371:661–664, 2010.

[26] M. D. Voisei and C. Zălinescu. Linear monotone subspaces of locally convex spaces. *Set-Valued Var. Anal.*, 18(1):29–55, 2010.

[27] M. D. Voisei and C. Zălinescu. Maximal monotonicity criteria for the composition and the sum under weak interiority conditions. *Math. Program.*, 123(1, Ser. B):265–283, 2010.

[28] C. Zălinescu. *Convex analysis in general vector spaces*. World Scientific Publishing Co. Inc., River Edge, NJ, 2002.